

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:14:14 ; Search time 157 Seconds
 (without alignments)
 282.548 Million cell updates/sec

Title: US-10-057-890A-31
 Perfect score: 884
 Sequence: 1 MKVSVAALSCMLMLVTALGSM.....GLNNCSSSNRLDGHQRVHAA 157

Scoring table: BLOSUM62
 Gapop 10.0 , Gapext 0.5

Searched: 1586107 seqs, 282547505 residues

Total number of hits satisfying chosen parameters: 1586107

Minimum DB seq length: 0
 Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
 Maximum Match 100%
 Listing first 45 summaries

Database : A_Geneseq_29Jan04:*
 1: geneseqp1980s:*
 2: geneseqp1990s:*
 3: geneseqp2000s:*
 4: geneseqp2001s:*
 5: geneseqp2002s:*
 6: geneseqp2003as:*
 7: geneseqp2003bs:*
 8: geneseqp2004s:*

Pred. No. is the number of results predicted by chance to have a
 score greater than or equal to the score of the result being printed,
 and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	% Query		DB	ID	Description
		Match	Length			
1	884	100.0	157	5	ABG32540	Abg32540 Human CCR
2	797	90.2	138	5	ABG32539	Abg32539 Human CCR
3	380	43.0	439	2	AAy41280	Aay41280 Fusion pr
4	379	42.9	352	2	AAW27407	Aaw27407 Human CCR
5	379	42.9	352	2	AAW27123	Aaw27123 Human che
6	379	42.9	352	2	AAW23835	Aaw23835 Human CC
7	379	42.9	352	2	AAW88232	Aaw88232 HIV-1 co-
8	379	42.9	352	4	AAE07048	Aae07048 Human G-p
9	379	42.9	352	4	AAG80111	Aag80111 Human CCR

10	379	42.9	352	4	AAE04321	Aae04321	Human	che
11	379	42.9	352	4	AAE07039	Aae07039	Human	G-p
12	379	42.9	352	4	AAB46858	Aab46858	Human	HDG
13	379	42.9	352	4	ABB56342	Abb56342	Non-endog	
14	379	42.9	352	4	AAB83354	Aab83354	Human	CCR
15	379	42.9	352	4	AAB82948	Aab82948	Human	HIV
16	379	42.9	352	5	AAU97152	Aau97152	Human	G-p
17	379	42.9	352	5	AAM52829	Aam52829	Human	CCR
18	379	42.9	352	5	AAM52828	Aam52828	Human	CC
19	379	42.9	352	5	ABG70597	Abg70597	Human	G-p
20	379	42.9	352	5	ABG92883	Abg92883	Human	imm
21	379	42.9	352	5	AAE25811	Aae25811	Human	G-p
22	379	42.9	352	5	ABB81054	Abb81054	G-protein	
23	379	42.9	352	5	ABB08343	Abb08343	Human	che
24	379	42.9	352	6	ABG75540	Abg75540	Human	G-p
25	379	42.9	352	6	ABR58602	Abr58602	Human	can
26	379	42.9	352	6	AAO29514	Aao29514	Human	C-C
27	379	42.9	352	6	ABU61654	Abu61654	Human	G-p
28	379	42.9	352	6	ABP97728	Abp97728	Amino aci	
29	379	42.9	352	6	ABP81933	Abp81933	Human	C-C
30	379	42.9	352	7	ADC03341	Adc03341	Human	che
31	374	42.3	371	2	AAW23834	Aaw23834	Human	CC
32	373	42.2	352	2	AAW07602	Aaw07602	Human	G-p
33	373	42.2	352	3	AAV80128	Aay80128	Human	G-p
34	373	42.2	352	4	AAE07046	Aae07046	Human	G-p
35	373	42.2	352	4	AAE07037	Aae07037	Human	G-p
36	373	42.2	352	5	AAU97150	Aau97150	Human	G-p
37	373	42.2	352	5	ABG92880	Abg92880	Human	G-p
38	373	42.2	352	5	AAE25808	Aae25808	Human	G-p
39	363	41.1	352	2	AAW27125	Aaw27125	Macaque c	
40	363	41.1	352	7	ADC03359	Adc03359	Macaque c	
41	359	40.6	352	4	AAG79089	Aag79089	Amino aci	
42	272	30.8	354	2	AAW54037	Aaw54037	Mouse	CC-
43	258	29.2	184	2	AAW27406	Aaw27406	Inactive	
44	258	29.2	215	2	AAW27408	Aaw27408	Inactive	
45	258	29.2	215	2	AAW88238	Aaw88238	HIV-1 co-	

ALIGNMENTS

RESULT 1

ABG32540

ID ABG32540 standard; protein; 157 AA.

XX

AC ABG32540;

XX

DT 15-NOV-2002 (first entry)

XX

DE Human CCR5-based scaffolded fusion protein #2.

XX

KW Scaffolded protein; CCR5; HIV; human immunodeficiency virus infection;

KW ECD; extracellular domain; metal chelating motif; zinc finger protein;

KW integral membrane protein; soluble loop; intracellular domain; ICD;

KW gene therapy; immunogen; viral infection; human.

XX

OS Homo sapiens.

OS Synthetic.

XX

FH Key Location/Qualifiers

FT Peptide 1. .19

FT /label= Signal_peptide

FT Protein 20. .157

FT /label= Mature_scaffolded_protein

XX

PN WO200260477-A1.

XX

PD 08-AUG-2002.

XX

PF 29-JAN-2002; 2002WO-US002377.

XX

PR 31-JAN-2001; 2001US-0265782P.

PR 31-JAN-2001; 2001US-0265858P.

XX

PA (HUMA-) HUMAN GENOME SCI INC.

XX

PI Coleman TA, Mansfield B;

XX

DR WPI; 2002-643357/69.

XX

PT Novel scaffolded fusion polypeptide useful for therapeutic purposes or
PT for screening molecules that bind/activate/inhibit/modulate the
PT polypeptide, comprises a functional polypeptide domain fused to a
PT scaffold domain.

XX

PS Example 2; Page 41; 64pp; English.

XX

CC The invention relates to a scaffolded fusion polypeptide comprising a
CC functional polypeptide domain fused to a scaffold domain, where the
CC functional polypeptide domain corresponds to a soluble loop of an
CC integral membrane protein (e.g. human CCR5, a transmembrane receptor
CC involved in HIV (human immunodeficiency virus) infection). Also included
CC are; (1) a polypeptide comprising a scaffold domain; (2) a nucleic acid
CC encoding the fusion polypeptide; (3) a vector cassette for the expression
CC of the fusion polypeptide comprising an expression region operably linked
CC to a promoter, where the expression region comprises a number of
CC cassettes, each of which encodes a module, domain or strand of the fusion
CC polypeptide and (4) a host cell comprising the vector or nucleic acid.
CC The fusion polypeptide is useful for screening molecules that
CC bind/activate/inhibit/modulate the fusion polypeptide, by expressing the
CC fusion polypeptide from and identifying a molecule that binds to the
CC fusion polypeptide. The fusion polypeptide is useful in diagnostic
CC methods, in assays to identify compounds that interact with loops of
CC fragments of an extracellular domain (ECD) or an intracellular domain
CC (ICD) or to rapidly assay the function of mutated portions of mutant
CC integral membrane proteins without having to produce significant
CC quantities of the entire mutant integral membrane protein, to generate
CC antibodies that recognise the integral membrane proteins from which they
CC are designed, to competitively bind the ligand of a naturally occurring
CC receptor in vitro or in vivo, to display and/or screen soluble domains
CC from protein such as integral membrane proteins, to probe the structure
CC of ECD or ICD, or both, of an integral protein membrane, to modulate the
CC activity of a receptor in vivo, and for treating or preventing viral
CC infection, preferably human HIV infection e.g. by gene therapy using the

CC encoding nucleic acid. The present sequence is a scaffolded protein based
CC on the ECD region of human CCR5 (not defined)
XX
SQ Sequence 157 AA;

Query Match 100.0%; Score 884; DB 5; Length 157;
Best Local Similarity 100.0%; Pred. No. 5.1e-84;
Matches 157; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

```
Qy      1 MKVSVAALSCLMLVTALGSMQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAAQW 60
          |||
Db      1 MKVSVAALSCLMLVTALGSMQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAAQW 60

Qy     61 DFGNTMCQHQRVHGHHHSYKCGLCSTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQR 120
          |||
Db     61 DFGNTMCQHQRVHGHHHSYKCGLCSTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQR 120

Qy    121 VHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVHAA 157
          |||
Db    121 VHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVHAA 157
```

RESULT 2

ABG32539

ID ABG32539 standard; protein; 138 AA.

XX

AC ABG32539;

XX

DT 15-NOV-2002 (first entry)

XX

DE Human CCR5-based scaffolded fusion protein #1.

XX

KW Scaffolded protein; CCR5; HIV; human immunodeficiency virus infection;
KW ECD; extracellular domain; metal chelating motif; zinc finger protein;
KW integral membrane protein; soluble loop; intracellular domain; ICD;
KW gene therapy; immunogen; viral infection; human.

XX

OS Homo sapiens.

OS Synthetic.

XX

PN WO200260477-A1.

XX

PD 08-AUG-2002.

XX

PF 29-JAN-2002; 2002WO-US002377.

XX

PR 31-JAN-2001; 2001US-0265782P.

PR 31-JAN-2001; 2001US-0265858P.

XX

PA (HUMA-) HUMAN GENOME SCI INC.

XX

PI Coleman TA, Mansfield B;

XX

DR WPI; 2002-643357/69.

XX

PT Novel scaffolded fusion polypeptide useful for therapeutic purposes or
PT for screening molecules that bind/activate/inhibit/modulate the

PT polypeptide, comprises a functional polypeptide domain fused to a
PT scaffold domain.

XX

PS Example 1; Page 21; 64pp; English.

XX

CC The invention relates to a scaffolded fusion polypeptide comprising a
CC functional polypeptide domain fused to a scaffold domain, where the
CC functional polypeptide domain corresponds to a soluble loop of an
CC integral membrane protein (e.g. human CCR5, a transmembrane receptor
CC involved in HIV (human immunodeficiency virus) infection). Also included
CC are; (1) a polypeptide comprising a scaffold domain; (2) a nucleic acid
CC encoding the fusion polypeptide; (3) a vector cassette for the expression
CC of the fusion polypeptide comprising an expression region operably linked
CC to a promoter, where the expression region comprises a number of
CC cassettes, each of which encodes a module, domain or strand of the fusion
CC polypeptide and (4) a host cell comprising the vector or nucleic acid.
CC The fusion polypeptide is useful for screening molecules that
CC bind/activate/inhibit/modulate the fusion polypeptide, by expressing the
CC fusion polypeptide from and identifying a molecule that binds to the
CC fusion polypeptide. The fusion polypeptide is useful in diagnostic
CC methods, in assays to identify compounds that interact with loops of
CC fragments of an extracellular domain (ECD) or an intracellular domain
CC (ICD) or to rapidly assay the function of mutated portions of mutant
CC integral membrane proteins without having to produce significant
CC quantities of the entire mutant integral membrane protein, to generate
CC antibodies that recognise the integral membrane proteins from which they
CC are designed, to competitively bind the ligand of a naturally occurring
CC receptor in vitro or in vivo, to display and/or screen soluble domains
CC from protein such as integral membrane proteins, to probe the structure
CC of ECD or ICD, or both, of an integral protein membrane, to modulate the
CC activity of a receptor in vivo, and for treating or preventing viral
CC infection, preferably human HIV infection e.g. by gene therapy using the
CC encoding nucleic acid. The present sequence is a scaffolded protein based
CC on the ECD region of human CCR5 (not defined)

XX

SQ Sequence 138 AA;

Query Match 90.2%; Score 797; DB 5; Length 138;
Best Local Similarity 100.0%; Pred. No. 5.1e-75;
Matches 138; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy	20	MDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAQWDFGNTMCQHQRVHGHHS	79
Db	1	MDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAQWDFGNTMCQHQRVHGHHS	60
Qy	80	YKGLCTRSQKEGLHYTCSSHFYPYSQYQFWKNFQTLKIHQRVHGGGGSYKGLCQEFFGL	139
Db	61	YKGLCTRSQKEGLHYTCSSHFYPYSQYQFWKNFQTLKIHQRVHGGGGSYKGLCQEFFGL	120
Qy	140	NNCSSNRLDGHQRVHAA	157
Db	121	NNCSSNRLDGHQRVHAA	138

RESULT 3

AAY41280

ID AAY41280 standard; protein; 439 AA.

XX
 AC AAY41280;
 XX
 DT 11-FEB-2000 (first entry)
 XX
 DE Fusion protein containing human chemokine receptor CCR-5.
 XX
 KW Prostaglandin; PG; E2EP3 receptor; E2EP2 receptor; CCR-5; human;
 KW chemokine receptor; ss2 adrenergic receptor; small G-protein rho;
 KW renal outer medullary potassium ion channel protein; ion-channel protein;
 KW lambda phage repressor protein; G-protein coupled receptor; bacteria;
 KW biochemical; vaccine; immunohistochemical; orphan receptor; HIV.
 XX
 OS Synthetic.
 OS Homo sapiens.
 XX
 PN WO9953033-A1.
 XX
 PD 21-OCT-1999.
 XX
 PF 16-APR-1999; 99WO-US008214.
 XX
 PR 16-APR-1998; 98US-0081989P.
 XX
 PA (UYVA-) UNIV VANDERBILT.
 XX
 PI Breyer RM, Ma L, Kennedy C;
 XX
 DR WPI; 1999-620416/53.
 DR N-PSDB; AAZ24738.
 XX
 PT New nucleic acid constructs for high level expression of eukaryotic
 PT proteins in bacteria, for producing e.g. chemokine receptor CCR-5 for
 PT preventing HIV infection.
 XX
 PS Disclosure; Page 49-53; 81pp; English.
 XX
 CC The invention provides isolated nucleic acid sequences that encode rabbit
 CC prostaglandin (PG) E2EP3 receptor, human PG E2EP2 receptor, human
 CC chemokine receptor CCR-5, human ss2 adrenergic receptor, rat renal outer
 CC medullary potassium ion channel protein or human small G-protein rho,
 CC together with deduced protein sequences. Also provided is a method for
 CC the production of eukaryotic proteins by culturing bacteria transformed
 CC with vectors containing the above nucleic acid sequences or a nucleic
 CC acid (I) that comprises: (i) first sequence that encodes either a
 CC sequence comprising at least three positively charged amino acids, or a
 CC DNA-binding protein, or a lambda phage repressor protein, placed upstream
 CC of, and in frame with, (ii) a sequence encoding a protein. (I) are used
 CC for recombinant production of eukaryotic proteins, particularly membrane
 CC proteins, G-protein coupled receptors or ion-channel proteins, in
 CC bacteria. These proteins are useful for biochemical or structural studies
 CC ; as therapeutic agents; in diagnostic and screening assays and as
 CC antigens for use in vaccines, and for raising antibodies that are useful
 CC as immunohistochemical markers, e.g. for orphan receptors or ion
 CC channels. Antibodies raised against the chemokine receptor CCR-5 can be
 CC used (when administered as antiserum or generated in vivo) to prevent
 CC entry of human immune deficiency virus (HIV) into cells

XX

SQ Sequence 439 AA;

Query Match 43.0%; Score 380; DB 2; Length 439;

Best Local Similarity 34.1%; Pred. No. 5.9e-31;

Matches 103; Conservative 8; Mismatches 31; Indels 160; Gaps 8;

```
Qy      1 MKVSVAALSCLMLVTALGSMQVSSPIYDINYTTSEPCQKINVKQIAA----- 49
          :|||||                                     |||||
Db      70 LKVSVEEF-----HMDYQVSSPIYDINYTTSEPCQKINVKQIAARLLPPLYSLVF 119

Qy      50 -----YKCGLC-----AAAQWDFGNTMC 67
          | |                                     |||||
Db     120 IFGFVGNMLVILILINCKRLKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC 179

Qy      68 QHQRVHGHHSYKCG----- 83
          | : | : |
Db     180 --QLLTGLYFIGFFSGIFFIILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFA 237

Qy      84 -----LCTRSQKEGLHYTCSSHFYPYSQYQFWKNFQTLKI----- 117
          : |||||
Db     238 SLPGIIFTRSQKEGLHYTCSSHFYPYSQYQFWKNFQTLKIVILGLVLPLLVMVICYSGILK 297

Qy     118 -----HQRVHGG-----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQ 152
          :: | | |||||
Db     298 TLLRCRNEKKRHRAVRLIFTIMIVYFLFWAPYNIVLLNLTFFQEFFGLNNCSSSNRLDQAM 357

Qy     153 RV 154
          :|
Db     358 QV 359
```

RESULT 4

AAW27407

ID AAW27407 standard; protein; 352 AA.

XX

AC AAW27407;

XX

DT 14-APR-1998 (first entry)

XX

DE Human CCR5.

XX

KW Human Cys-Cys chemokine receptor 5; CCR5; human immunodeficiency virus;
KW type 1; type 2; HIV-1; HIV-2; diagnosis; treatment; prevention;
KW inflammatory disease; rheumatoid arthritis; glomerulonephritis; asthma;
KW idiopathic pulmonary fibrosis; psoriasis; viral infection; cancer;
KW atherosclerosis; autoimmune disorder.

XX

OS Homo sapiens.

XX

PN WO9732019-A2.

XX

PD 04-SEP-1997.

XX

PF 28-FEB-1997; 97WO-BE000023.

XX

PR 01-MAR-1996; 96EP-00870021.

PR 06-AUG-1996; 96EP-00870102.
 XX
 PA (EURO-) EUROSCREEN SA.
 XX
 PI Samson M, Parmentier M, Vassart G, Libert F;
 XX
 DR WPI; 1997-479829/44.
 DR N-PSDB; AAT90117.
 XX
 PT Active and inactive forms of human CC chemokine receptor CCR-5 - useful
 PT to diagnose, prevent and/or treat inflammatory disorders, autoimmune
 PT disease and viral infection.
 XX
 PS Claim 4; Fig 1b-c; 94pp; English.
 XX
 CC The present sequence is human CC (Cys-Cys) chemokine receptor 5 (CCR5),
 CC which is stimulated by MIP-1 alpha, MIP-1 beta or RANTES chemokines, but
 CC not by monocyte chemoattractant protein 1 (MCP-1), MCP-2, MCP-3,
 CC interleukin-8 (IL-8) or growth related gene product alpha (GRO alpha)
 CC chemokines. Active CCR-5 is also a receptor of human immunodeficiency
 CC virus type 1 or type 2 (HIV-1 or HIV-2). CCR5 or its cDNA can used to
 CC diagnose, treat and/or prevent inflammatory diseases, e.g. rheumatoid
 CC arthritis, glomerulonephritis, asthma, idiopathic pulmonary fibrosis and
 CC psoriasis, viral infections, especially HIV-1 or HIV-2 infection, cancer,
 CC atherosclerosis and autoimmune disorders
 XX
 SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 2; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||||||||||||||              :: |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC---QEFGFLNNCSSSNRLDGHQRV 154
          |  |                      |||||  | : |
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 5
 AAW27123
 ID AAW27123 standard; protein; 352 AA.
 XX

AC AAW27123;
 XX
 DT 14-DEC-1997 (first entry)
 XX
 DE Human chemokine receptor 88C.
 XX
 KW Chemokine receptor 88C; atherosclerosis; rheumatoid arthritis; tumour;
 KW asthma; viral infection; AIDS; inflammation; autoimmune disease; therapy;
 KW diagnosis; leukocyte trafficking; G protein coupled receptor; ligand;
 KW modulator; antibody; human.
 XX
 OS Homo sapiens.
 XX
 FH Key Location/Qualifiers
 FT Domain 1. .32
 FT /label= Extracellular_domain
 FT Domain 56. .67
 FT /label= Intracellular_domain
 FT Domain 89. .112
 FT /label= Extracellular_domain
 FT Domain 125. .145
 FT /label= Intracellular_domain
 FT Domain 166. .191
 FT /label= Extracellular_domain
 FT Domain 213. .235
 FT /label= Intracellular_domain
 FT Domain 259. .280
 FT /label= Extracellular_domain
 FT Domain 301. .352
 FT /label= Intracellular_domain
 XX
 PN WO9722698-A2.
 XX
 PD 26-JUN-1997.
 XX
 PF 20-DEC-1996; 96WO-US020759.
 XX
 PR 20-DEC-1995; 95US-00575967.
 PR 07-JUN-1996; 96US-00661393.
 XX
 PA (ICOS-) ICOS CORP.
 XX
 PI Gray PW, Schweickart VL, Raport CJ;
 XX
 DR WPI; 1997-341689/31.
 DR N-PSDB; AAT85161.
 XX
 PT New nucleic acid encoding chemokine receptors 88-2B and 88C - used to
 PT modulate leukocyte trafficking, e.g. for treatment of inflammation,
 PT tumours, viral infections, auto-immune diseases, etc.
 XX
 PS Claim 16; Page 47-48; 65pp; English.
 XX
 CC This polypeptide sequence comprises novel human chemokine receptor 88C, a
 CC G protein coupled receptor that is involved in leukocyte trafficking. Its
 CC amino sequence was deduced from a cDNA clone (AAT85161) isolated from a
 CC macrophage library. It shows 62% identity to CCCKR1. Chemokine receptor

CC 88-2B (see AAW27124) has also been identified. 88C and 88-2B receptors
 CC and their polypeptide fragments can be produced in transformed host
 CC cells. The receptors, peptides comprising one or more of the
 CC extracellular or intracellular domains, and anti-receptor antibodies can
 CC be used to modulate receptor activities, particularly ligand and G
 CC protein binding, and are potentially potentially useful in the treatment
 CC of atherosclerosis, rheumatoid arthritis, tumours, asthma, viral
 CC infection, AIDS, inflammatory conditions, pathological immune response,
 CC abnormal haematopoietic processes etc
 XX
 SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 2; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |
Db      61 LKSMTDIYLLNLAIISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  |
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

RESULT 6

AAW23835

ID AAW23835 standard; protein; 352 AA.

XX

AC AAW23835;

XX

DT 08-JUN-1998 (first entry)

XX

DE Human CC chemokine receptor 5 (CCR5).

XX

KW CC chemokine receptor 5; CCR5; G-protein coupled receptor;

KW human immunodeficiency virus; HIV; CD4; AIDS; therapy; transgenic animal.

XX

OS Homo sapiens.

XX

FH Key Location/Qualifiers

FT Domain 29. .55

FT /label= I

FT /note= "transmembrane domain"

FT Domain 104. .126

FT /label= III
 FT /note= "transmembrane domain"
 FT Region 109. .120
 FT /note= "extracellular loop-1 (Claim 19)"
 FT Domain 143. .171
 FT /label= IV
 FT /note= "transmembrane domain"
 FT Region 187. .210
 FT /note= "extracellular loop-2 (Claim 19)"
 FT Domain 194. .219
 FT /label= V
 FT /note= "transmembrane domain"
 FT Domain 238. .258
 FT /label= VI
 FT /note= "transmembrane domain"
 FT Region 261. .276
 FT /note= "extracellular loop-3 (Claim 19)"
 FT Domain 277. .300
 FT /label= VII
 FT /note= "transmembrane domain"

XX

PN WO9745543-A2.

XX

PD 04-DEC-1997.

XX

PF 28-MAY-1997; 97WO-US009586.

XX

PR 28-MAY-1996; 96US-0018508P.

XX

PA (USSH) US DEPT HEALTH & HUMAN SERVICES.

XX

PI Combadiere C, Feng Y, Berger EA, Alkhatib G, Murphy PM;

PI Broder CC, Kennedy PE;

XX

DR WPI; 1998-032650/03.

DR

N-PSDB; AAT76920.

XX

PT CC chemokine receptor 5 polypeptide - used to inhibit membrane fusion

PT

XX

PS

XX

CC This protein sequence comprises of a novel human macrophage-selective CC
 CC chemokine receptor that has been designated CCR5. The sequence was
 CC deduced from an isolated cDNA clone (see AAT76920). An Ala127Leu variant
 CC (see W238340 of CCR5 was also identified. The susceptibility of human
 CC macrophages to HIV infection depends on cell surface expression of CD4
 CC and CCR5. CCR5 is a member of the 7-transmembrane superfamily of G-
 CC protein coupled cell surface molecules. It plays an essential role in the
 CC membrane fusion step of infection by some HIV isolates. The establishment
 CC of stable, non-human cell lines and transgenic mammals having cells that
 CC coexpress human CD4 and CCR5 provides valuable tools for research of HIV
 CC infection. Antibodies that bind to CCR5, CCR5 variants, and CCR5-binding
 CC agents capable of blocking membrane fusion between HIV and target cells
 CC represent potential anti-HIV therapeutics for macrophage tropic strains
 CC of HIV

XX

SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 2; Length 352;
Best Local Similarity 35.0%; Pred. No. 5.7e-31;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||  :: |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  |  |||||||||||  :|
Db     239 TIMIVYFLFWAPYNIVLLLNLTQEFFGLNNCSSSNRLDQAMQV 281
```

RESULT 7

AAW88232

ID AAW88232 standard; protein; 352 AA.

XX

AC AAW88232;

XX

DT 15-MAR-1999 (first entry)

XX

DE HIV-1 co-receptor CCR5.

XX

KW HIV-1; CCR5; CCR5m303; co-receptor; infection; diagnosis; AIDS;
KW gene therapy; human.

XX

OS Homo sapiens.

XX

FH Key Location/Qualifiers

FT Domain 32. .56

FT /note= "transmembrane domain 1"

FT Domain 67. .87

FT /note= "transmembrane domain 2"

FT Misc-difference 101

FT /note= "corresponds to TGT (Cys) in wild-type CCR5, TGA
(Stop) in CCR5m303"

FT Domain 103. .124

FT /note= "transmembrane domain 3"

FT Domain 142. .167

FT /note= "transmembrane domain 4"

FT Domain 200. .223

FT /note= "transmembrane domain 5"

FT Domain 236. .260


```

Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178
Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238
          |||
Qy      125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154
          | | |||
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281
          | | |||

```

RESULT 8

AAE07048

ID AAE07048 standard; protein; 352 AA.

XX

AC AAE07048;

XX

DT 16-OCT-2001 (first entry)

XX

DE Human G-protein chemokine receptor (CCR5) HDGNR10 protein #2.

XX

KW Human; G-protein chemokine receptor; CCR5; HDGNR10; inflammation; HIV;
 KW human immunodeficiency virus; antimicrobial; vasodilator; vulnerary;
 KW cytostatic; immunosuppressive; nootropic; neuroprotective; gene therapy;
 KW neurodegenerative disorder; Kaposi's sarcoma; autoimmune disease;
 KW rheumatoid arthritis; cancer; breast; ovary; adrenal gland; bone marrow;
 KW gastrointestinal tract; lung; liver; immune disorder; Addison's disease;
 KW haemolytic anaemia; autoimmune thyroiditis; diabetes mellitus; allergy;
 KW multiple sclerosis; ulcerative colitis; Crohn's disease; wound healing;
 KW cardiovascular disorder; myocardial ischaemia.

XX

OS Homo sapiens.

XX

PN WO200158916-A2.

XX

PD 16-AUG-2001.

XX

PF 09-FEB-2001; 2001WO-US004153.

XX

PR 09-FEB-2000; 2000US-0181258P.

PR 09-MAR-2000; 2000US-0187999P.

PR 22-SEP-2000; 2000US-0234336P.

XX

PA (HUMA-) HUMAN GENOME SCI INC.

XX

PI Rosen CA, Roschke V, Li Y, Ruben SM;

XX

DR WPI; 2001-488966/53.

DR N-PSDB; AAD13299.

XX

PT Isolated nucleic acid encoding a human G-protein chemokine receptor
 PT (CCR5) HDGNR10 polypeptide, useful for preventing or treating autoimmune
 PT diseases e.g. rheumatoid arthritis, hyperproliferative disorders and
 PT neurodegenerative disorders.

XX

PS Example 40; Page 504-505; 518pp; English.

XX

CC The invention relates to human G-protein chemokine receptor (CCR5)
 CC HDGNR10 polypeptides and polynucleotides. CCR5 HDGNR10 antibodies are
 CC useful for treating, preventing or ameliorating a disease or disorder
 CC associated with inflammation, defective or aberrant chemotaxis of immune
 CC cells, HIV infection (such as Pneumocystis carinii pneumonia or Kaposi's
 CC sarcoma) or defective or aberrant T-cell antigen presenting cell
 CC interaction. The disease or disorder may also be an infectious disease
 CC (e.g. a viral infection such as an early stage HIV infection, a
 CC cytomegalovirus infection, or a poxvirus infection), an autoimmune
 CC disease (e.g. rheumatoid arthritis) or a neurodegenerative disorder. The
 CC disease or disorder may be associated with aberrant CCR5 expression, lack
 CC of CCR5 function, aberrant CCR5 ligand expression, or lack of CCR5 ligand
 CC function. CCR5 HDGNR10 protein is used as a food additive or preservative
 CC to increase or decrease storage capabilities. CCR5 HDGNR10 DNA are useful
 CC for chromosome identification and in gene therapy. CCR5 HDGNR10 DNA,
 CC protein, antibodies, agonists and antagonists are also useful in the
 CC diagnosis, treatment and prevention of cancer (breast, ovary, adrenal
 CC gland, bone, bone marrow, gastrointestinal tract, liver, lung,
 CC urogenital); immune disorders (Addison's disease, allergies, autoimmune
 CC haemolytic anaemia, autoimmune thyroiditis, diabetes mellitus, Crohn's
 CC disease, multiple sclerosis, rheumatoid arthritis and ulcerative colitis)
 CC ; cardiovascular disorders (myocardial ischaemias) and wound healing. The
 CC present sequence is human CCR5 HDGNR10 protein
 XX
 SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          | |               |||
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC---QEFFGLNNCSSSNRLDGHQRV 154
          | |               |||
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 9

AAG80111

ID AAG80111 standard; protein; 352 AA.

XX

AC AAG80111;

XX

DT 17-JAN-2002 (first entry)
 XX
 DE Human CCR5 protein.
 XX
 KW Chemokine; tumour diagnosis; colorectal; prostatic; organ rejection;
 KW inflammation; autoimmune disease; metastasis; bronchial asthma; lupus;
 KW chronic bowel inflammation; rheumatoid arthritis; cytostatic;
 KW antiinflammatory; antiasthmatic; immunosuppressive; dermatological;
 KW antirheumatic; antiarthritic.
 XX
 OS Homo sapiens.
 XX
 PN WO200172830-A2.
 XX
 PD 04-OCT-2001.
 XX
 PF 02-APR-2001; 2001WO-EP003708.
 XX
 PR 31-MAR-2000; 2000DE-01016013.
 XX
 PA (IPFP-) IPF PHARM GMBH.
 PA (FORS/) FORSSMANN U.
 XX
 PI Forssmann W, Adermann K, Heitland A, Spodsberg N;
 XX
 DR WPI; 2001-626256/72.
 XX
 PT Diagnostic agent containing two or more receptor-specific ligands, useful
 PT for detecting tumors, inflammation etc., also therapeutic use of ligand
 PT inhibitors.
 XX
 PS Disclosure; Page 10; 26pp; German.
 XX
 CC This invention describes a novel diagnostic agent (A) comprising at least
 CC two different ligands (I) for receptors (II) that are implicated in
 CC disease. (A) are used for the diagnosis of tumors (especially colorectal
 CC or prostatic), organ rejection, inflammation and autoimmune diseases.
 CC Also inhibitors of (I) are used therapeutically against tumors (and their
 CC metastases), inflammation (particularly bronchial asthma or chronic bowel
 CC inflammation), or autoimmune diseases (rheumatoid arthritis or lupus),
 CC where the (cardio)vascular, lymphatic, respiratory, nervous, digestive,
 CC endocrine, motor or urogenital systems or skin are affected, and bone
 CC marrow diseases. The products of the invention are chemokine derivatives
 CC which have cytostatic, antiinflammatory, antiasthmatic,
 CC immunosuppressive, dermatological, antirheumatic, antiarthritic.
 CC Chemokines act on specific tumor and inflammatory cells through a
 CC constellation of chemokine receptors (CR), which control migration and
 CC proliferation of these cells. AAG80045-AAG80128 represent human chemokine
 CC fragments used to illustrate the method of the invention
 XX
 SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49


```

      |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
QY      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
      | | ||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
QY      84 -----LCTRSQKEGLHYTC 97
      : |||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178
QY      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      ||| : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238
QY      125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154
      | | ||| : |
Db      239 TIMIVYFLEWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281

```

RESULT 10

AAE04321

ID AAE04321 standard; protein; 352 AA.

XX

AC AAE04321;

XX

DT 04-SEP-2001 (first entry)

XX

DE Human chemokine receptor (CKR), CC-CKR-5 related protein #2.

XX

KW Human; transformed mammalian cell; CD4; reporter gene; translocation;
 KW human immuno deficiency virus; HIV; long terminal repeat; LTR; therapy;
 KW chemokine receptor; CKR; cellular dysfunction; HIV infection; cofactor;
 KW CC-CKR-5; envelope glycoprotein; anti-HIV.

XX

OS Homo sapiens.

XX

PN US6258527-B1.

XX

PD 10-JUL-2001.

XX

PF 21-MAY-1997; 97US-00861105.

XX

PR 20-MAY-1996; 96US-0017157P.

PR 19-JUN-1996; 96US-0020043P.

PR 19-MAY-1997; 97US-00858660.

XX

PA (AARO-) AARON DIAMOND AIDS RES CENT.

PA (UYN Y) UNIV NEW YORK STATE.

XX

PI Littman DR, Deng H, Ellmeier W, Landau NR, Liu R;

XX

DR WPI; 2001-417127/44.

DR N-PSDB; AAD08577.

XX

PT Transformed mammalian cell (I) that contains a CD4 gene, reporter gene
 PT and HIV LTR for identification of drugs and antibodies for treatment of

PT HIV.
 XX
 PS Disclosure; Col 47-50; 37pp; English.
 XX
 CC The present invention relates to a transformed mammalian cell that
 CC contains a gene encoding CD4, a construct encoding a reporter gene under
 CC the regulation of an human immuno deficiency virus (HIV) long terminal
 CC repeat (LTR) and that has been transduced with a vector encoding a human
 CC chemokine receptor (CKR) where the CD4 and the CKR are present on the
 CC cell surface of transformed mammalian cell. The invention is useful for
 CC identifying drugs or antibodies that interfere with the translocation of
 CC HIV into transformed mammalian cell or for identifying a human chemokine
 CC receptor that facilitates the infection of a particular HIV strain into
 CC the transformed mammalian cell. Compounds identified can be used to treat
 CC cellular dysfunction and to prevent or combat HIV infection. The present
 CC sequence is a human chemokine receptor (CKR), CC-CKR-5 related protein.
 CC CC-CKR-5 is the principal cofactor for entry mediated by the envelope
 CC glycoproteins of primary macrophage-tropic strains of HIV-1
 XX
 SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||||||||||||||||||||||||||
 Db 1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
 | | ||||||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||||||||
 Db 119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGLIIFTRSQKEGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 |||||||||||||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
 | | ||||||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLNLTQEFFGLNNCSSSNRLDQAMQV 281

RESULT 11

AAE07039

ID AAE07039 standard; protein; 352 AA.

XX

AC AAE07039;

XX

DT 16-OCT-2001 (first entry)

XX

DE Human G-protein chemokine receptor (CCR5) HDGNR10 protein #2.

XX

KW Human; G-protein chemokine receptor; CCR5; HDGNR10; inflammation; HIV;

KW human immunodeficiency virus; antimicrobial; vasodilator; vulnerary;
 KW cytostatic; immunosuppressive; nootropic; neuroprotective; gene therapy;
 KW neurodegenerative disorder; Kaposi's sarcoma; autoimmune disease;
 KW rheumatoid arthritis; cancer; breast; ovary; adrenal gland; bone marrow;
 KW gastrointestinal tract; lung; liver; immune disorder; Addison's disease;
 KW haemolytic anaemia; autoimmune thyroiditis; diabetes mellitus; allergy;
 KW multiple sclerosis; ulcerative colitis; Crohn's disease; wound healing;
 KW cardiovascular disorder; myocardial ischaemia.
 XX
 OS Homo sapiens.
 XX
 PN WO200158915-A2.
 XX
 PD 16-AUG-2001.
 XX
 PF 09-FEB-2001; 2001WO-US004152.
 XX
 PR 09-FEB-2000; 2000US-0181258P.
 PR 09-MAR-2000; 2000US-0187999P.
 PR 22-SEP-2000; 2000US-0234336P.
 XX
 PA (HUMA-) HUMAN GENOME SCI INC.
 XX
 PI Rosen CA, Roschke V, Li Y, Ruben SM;
 XX
 DR WPI; 2001-488965/53.
 DR N-PSDB; AAD13198.
 XX
 PT Isolated nucleic acid encoding a human G-protein chemokine receptor
 PT (CCR5) HDGNR10 polypeptide, useful for preventing or treating autoimmune
 PT diseases e.g. rheumatoid arthritis, hyperproliferative disorders and
 PT neurodegenerative disorders.
 XX
 PS Example 40; Page 486-487; 495pp; English.
 XX
 CC The present sequence is human G-protein chemokine receptor (CCR5) HDGNR10
 CC protein. CCR5 HDGNR10 antibodies are useful for treating, preventing or
 CC ameliorating a disease or disorder associated with inflammation,
 CC defective or aberrant chemotaxis of immune cells, HIV infection (such as
 CC Pneumocystis carinii pneumonia or Kaposi's sarcoma) or defective or
 CC aberrant T-cell antigen presenting cell interaction. The disease or
 CC disorder may also be an infectious disease (e.g. a viral infection such
 CC as an early stage HIV infection, a cytomegalovirus infection, or a
 CC poxvirus infection), an autoimmune disease (e.g. rheumatoid arthritis) or
 CC a neurodegenerative disorder. The disease or disorder may be associated
 CC with aberrant CCR5 expression, lack of CCR5 function, aberrant CCR5
 CC ligand expression, or lack of CCR5 ligand function. CCR5 HDGNR10 protein
 CC is used as a food additive or preservative to increase or decrease
 CC storage capabilities. CCR5 HDGNR10 DNA are useful for chromosome
 CC identification and in gene therapy. CCR5 HDGNR10 DNA, protein,
 CC antibodies, agonists and antagonists are also useful in the diagnosis,
 CC treatment and prevention of cancer (breast, ovary, adrenal gland, bone,
 CC bone marrow, gastrointestinal tract, liver, lung, urogenital); immune
 CC disorders (Addison's disease, allergies, autoimmune haemolytic anaemia,
 CC autoimmune thyroiditis, diabetes mellitus, Crohn's disease, multiple
 CC sclerosis, rheumatoid arthritis and ulcerative colitis); cardiovascular
 CC disorders (myocardial ischaemias) and wound healing

XX

SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;

Best Local Similarity 35.0%; Pred. No. 5.7e-31;

Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||||  :: |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYS GILKTLLRCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC---QEFGNLCSSSNRLDGHQRV 154
          |  |  |||||||||||||  :|
Db     239 TIMIVYFLEWAPYNIVLLLNTFQEFGNLCSSSNRLDQAMQV 281
```

RESULT 12

AAB46858

ID AAB46858 standard; protein; 352 AA.

XX

AC AAB46858;

XX

DT 16-AUG-2001 (revised)

DT 02-AUG-2001 (revised)

DT 04-MAY-2001 (first entry)

XX

DE Human HDG NR10 protein.

XX

KW HDG NR10; human; G-protein chemokine receptor; antiinflammatory;
KW immunomodulatory; anticoagulant; antiallergic; immunosuppressive;
KW cytostatic; antiparasitic; antipsoriatic; antirheumatic; antiarthritic;
KW vasotropic; gene therapy; haematopoiesis; wound healing; coagulation;
KW angiogenesis; solid tumour; infection; leukemia; growth factor activity;
KW T-cell mediated autoimmune disease; psoriasis; allergy; atherogenesis;
KW anaphylaxis; malignancy; inflammation; histamine; IgE; silicosis; shock;
KW immunoglobulin E-mediated allergic reaction; rheumatoid arthritis;
KW prostaglandin-independent fever; bone marrow failure; sarcoidosis;
KW hyper-eosinophilic syndrome; vulnerary.

XX

OS Homo sapiens.

XX

PN US2001000241-A1.

XX

PD 12-APR-2001.

XX

PF 29-NOV-2000; 2000US-00725285.

XX

PR 06-JUN-1995; 95US-00466343.

PR 18-NOV-1998; 98US-00195662.

PR 25-JUN-1999; 99US-00339912.

XX

PA (LIYY/) LI Y.

PA (RUBE/) RUBEN S M.

XX

PI Li Y, Ruben SM;

XX

DR WPI; 2001-226317/23.

DR N-PSDB; AAF26390.

XX

PT New human G-protein chemokine receptor polypeptides and polynucleotides,
PT useful for identifying (ant)agonists to the G-protein chemokine receptor.

XX

PS Claim 1a; Page 15; 22pp; English.

XX

CC This invention describes a novel receptor polypeptide (I) selected from
CC (i) a fully defined 329 amino acid sequence (II) fully disclosed in the
CC specification; and (ii) a polypeptide encoded by the cDNA contained in a
CC plasmid, and fragments, analogs and derivatives of the polypeptide. The
CC products of the invention have antiinflammatory, immunomodulatory,
CC anticoagulant, antiallergic, immunosuppressive, vulnerary, cytostatic,
CC antiparasitic, antipsoriatic, antirheumatic, antiarthritic and vasotropic
CC activity and can be used for gene therapy. The G-protein chemokine
CC receptors, HDGNR10, (I) are useful for screening for compounds which
CC activate or inhibit activation of (I). The products of the invention can
CC also be used for stimulating haematopoiesis, wound healing, coagulation,
CC angiogenesis, treating solid tumours, chronic infections, leukemia, T-
CC cell mediated autoimmune diseases, parasitic infections, psoriasis, and
CC stimulating growth factor activity. HDGNR10 is useful for treating
CC allergy, atherogenesis, anaphylaxis, malignancy, chronic and acute
CC inflammation, histamine and immunoglobulin E (IgE)-mediated allergic
CC reactions, prostaglandin-independent fever, bone marrow failure,
CC silicosis, sarcoidosis, rheumatoid arthritis, shock and hyper-
CC eosinophilic syndrome. (N.B. This record was resubmitted to correct
CC errors in the keyword formatting)

XX

SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;

Best Local Similarity 35.0%; Pred. No. 5.7e-31;

Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      |||
Db 1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy 50 -----YKGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | ||| : | : |
Db 61 LKSMTDIYLLNLAISDLFFLLTVPEWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy 84 -----LCTRSQKEGLHYTC 97
      : |||
Db 119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178
```

```

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      |||||||||||||||||||
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLRCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
      | | ||||||||||| :|
Db      239 TIMIVYFLFWAPYNIVLLINTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 13

ABB56342

ID ABB56342 standard; protein; 352 AA.

XX

AC ABB56342;

XX

DT 18-FEB-2002 (first entry)

XX

DE Non-endogenous human GPCR protein, SEQ ID NO: 477.

XX

KW Human; G protein-coupled receptor; GPCR; non-endogenous; mutant;
 KW constitutively activated GPCR; agonist; disease.

XX

OS Homo sapiens.

OS Synthetic.

XX

PN WO200177172-A2.

XX

PD 18-OCT-2001.

XX

PF 05-APR-2001; 2001WO-US011098.

XX

PR 07-APR-2000; 2000US-0195747P.

XX

PA (AREN-) ARENA PHARM INC.

XX

PI Lehmann-Bruinsma K, Liaw CW, Lin I;

XX

DR WPI; 2001-648759/74.

DR

N-PSDB; ABI97978.

XX

PT Identifying agonists of G protein-coupled receptors (GPCRs) for use in
 PT disease treatment, comprises contacting candidate compounds with versions
 PT of GPCRs.

XX

PS Claim 1; Page 277-278; 394pp; English.

XX

CC The invention relates to G protein-coupled receptors (GPCRs) for which
 CC the endogenous ligand has been identified. Non-endogenous constitutively
 CC activated versions of known GPCRs are used in the invention for the
 CC direct identification of candidate compounds as receptor agonists,
 CC inverse agonists or partial agonists. Such agonists are useful as
 CC therapeutic agents for diseases or disorders associated with GPCRs. The
 CC present sequence is a non-endogenous version of a known human GPCR

XX

SQ Sequence 352 AA;

Query Match 42.9%; Score 379; DB 4; Length 352;
 Best Local Similarity 35.0%; Pred. No. 5.7e-31;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               |||
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRAKRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |   |               |||
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

RESULT 14

AAB83354

ID AAB83354 standard; protein; 352 AA.

XX

AC AAB83354;

XX

DT 09-OCT-2001 (first entry)

XX

DE Human CCR5 protein sequence.

XX

KW Chemotactic chemokine receptor 5; gp120; CD4; therapy; HIV; CCR5;
 KW human immunodeficiency virus; anti-inflammatory disease; human.

XX

OS Homo sapiens.

XX

PN EP1118858-A2.

XX

PD 25-JUL-2001.

XX

PF 03-JAN-2001; 2001EP-00300020.

XX

PR 12-JAN-2000; 2000GB-00000659.

PR 12-JAN-2000; 2000GB-00000661.

PR 12-JAN-2000; 2000GB-00000663.

XX

PA (PFIZ) PFIZER LTD.

PA (PFIZ) PFIZER INC.

XX

PI Dobbs S, Perros M, Rickett GA;

XX

DR WPI; 2001-477088/52.

DR N-PSDB; AAF87099.

XX
 DE Human HIV-1 co-receptor CCR5.
 XX
 KW CCR5; chemokine; co-receptor; human immunodeficiency virus type 1; HIV-1;
 KW infection; therapy; vaccine; anti-HIV-1.
 XX
 OS Homo sapiens.
 XX
 FH Key Location/Qualifiers
 FT Binding-site 2. .18
 FT /note= "binds to HIV-1 gp120"
 XX
 PN WO200164710-A2.
 XX
 PD 07-SEP-2001.
 XX
 PF 28-FEB-2001; 2001WO-US006699.
 XX
 PR 29-FEB-2000; 2000US-0185667P.
 PR 19-MAY-2000; 2000US-0205839P.
 PR 07-FEB-2001; 2001US-0267231P.
 XX
 PA (PROG-) PROGENICS PHARM INC.
 PA (AARO-) AARON DIAMOND AIDS RES CENT.
 XX
 PI Dragic T, Olson WC;
 XX
 DR WPI; 2001-611273/70.
 DR N-PSDB; AAH26903.
 XX
 PT Novel compounds comprising specific amino acids within CCR5 (HIV 1 co-
 PT receptor) amino terminal domain including negatively charged and two
 PT sulfated tyrosine residues is useful for treating HIV infection in
 PT humans.
 XX
 PS Claim 1; Page 30; 163pp; English.
 XX
 CC The present sequence is that of human HIV-1 co-receptor CCR5. Amino acids
 CC 2-18 in the N-terminal region of CCR5 comprise an HIV-1 gp120-binding
 CC site that determines the specificity of the interaction between CCR5 and
 CC HIV-1 gp120. Post-translational sulfation of the tyrosine residues in the
 CC CCR5 N-terminus is required for gp120 binding and may critically modulate
 CC the susceptibility of target cells to HIV-1 infection in vivo. The
 CC invention provides claimed sulfated peptides (see AAB82947) that are
 CC based on the CCR5 N-terminal region and which are effective for
 CC inhibiting HIV-1 binding to CCR5. These peptides are used in claimed
 CC methods of inhibiting HIV infection of CD4+ cells, of preventing CD4+
 CC cells from becoming infected with HIV, of treating a subject whose CD4+
 CC cells are infected with HIV, and of identifying an agent which inhibits
 CC binding of a CCR5 ligand to a CCR5 receptor. The methods may be carried
 CC out in a subject, especially a human, infected (therapeutic method), not
 CC infected with HIV (prophylactic method), or in a subject who is not
 CC infected with, but has been exposed to, HIV
 XX
 SQ Sequence 352 AA;

Query Match

42.9%; Score 379; DB 4; Length 352;

Best Local Similarity 35.0%; Pred. No. 5.7e-31;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               ||||||||| | : | : : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||||||||||||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC-----QEFFGLNNC SSSNRLDGHQRV 154
          |   |               ||||||||||||| :|
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281
```

Search completed: March 5, 2004, 16:22:51
Job time : 158 secs

GenCore version 5.1.6
Copyright (c) 1993 - 2004 Compugen Ltd.

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:17:14 ; Search time 42.1574 Seconds
(without alignments)
192.262 Million cell updates/sec

Title: US-10-057-890A-31
Perfect score: 884
Sequence: 1 MKVSVAALSCLMLVTALGSM.....GLNNCSSSNRLDGHQRVHAA 157

Scoring table: BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 389414 seqs, 51625971 residues

Total number of hits satisfying chosen parameters: 389414

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : Issued_Patents_AA:*
1: /cgn2_6/ptodata/2/iaa/5A_COMB.pep:*
2: /cgn2_6/ptodata/2/iaa/5B_COMB.pep:*
3: /cgn2_6/ptodata/2/iaa/6A_COMB.pep:*
4: /cgn2_6/ptodata/2/iaa/6B_COMB.pep:*
5: /cgn2_6/ptodata/2/iaa/PCTUS_COMB.pep:*
6: /cgn2_6/ptodata/2/iaa/backfiles1.pep:*

Pred. No. is the number of results predicted by chance to have a
score greater than or equal to the score of the result being printed,
and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	% Query		DB	ID	Description
		Match	Length			
1	379	42.9	352	3	US-09-087-232A-13	Sequence 13, Appl
2	379	42.9	352	3	US-08-861-105-14	Sequence 14, Appl
3	379	42.9	352	3	US-08-575-967A-2	Sequence 2, Appli
4	379	42.9	352	4	US-08-833-752-5	Sequence 5, Appli
5	379	42.9	352	4	US-09-502-783A-2	Sequence 2, Appli
6	379	42.9	352	4	US-09-796-202-1	Sequence 1, Appli
7	374	42.3	352	3	US-09-045-583-52	Sequence 52, Appl
8	374	42.3	352	4	US-09-534-185-52	Sequence 52, Appl
9	373	42.2	352	3	US-08-466-343D-2	Sequence 2, Appli
10	359	40.6	352	4	US-09-517-605-5	Sequence 5, Appli
11	272	30.8	354	4	US-08-724-984A-2	Sequence 2, Appli

12	258	29.2	184	4	US-08-833-752-4	Sequence 4, Appli
13	258	29.2	215	3	US-09-087-232A-17	Sequence 17, Appl
14	258	29.2	215	4	US-08-833-752-6	Sequence 6, Appli
15	182.5	20.6	100	3	US-09-087-232A-15	Sequence 15, Appl
16	159	18.0	87	3	US-09-087-232A-18	Sequence 18, Appl
17	148.5	16.8	488	2	US-08-933-750C-17	Sequence 17, Appl
18	148.5	16.8	488	3	US-09-234-613-17	Sequence 17, Appl
19	128	14.5	54	4	US-08-833-752-11	Sequence 11, Appl
20	128	14.5	313	4	US-09-800-729-196	Sequence 196, App
21	128	14.5	326	4	US-09-800-729-195	Sequence 195, App
22	123.5	14.0	717	4	US-09-881-578A-2	Sequence 2, Appli
23	123	13.9	22	3	US-08-861-105-4	Sequence 4, Appli
24	121	13.7	518	4	US-09-881-578A-4	Sequence 4, Appli
25	121	13.7	711	2	US-08-820-170A-10	Sequence 10, Appl
26	121	13.7	711	3	US-09-055-699-10	Sequence 10, Appl
27	121	13.7	711	3	US-09-273-565-10	Sequence 10, Appl
28	121	13.7	711	4	US-09-565-538-10	Sequence 10, Appl
29	121	13.7	711	4	US-09-661-468-10	Sequence 10, Appl
30	121	13.7	711	4	US-09-976-165-10	Sequence 10, Appl
31	120.5	13.6	185	4	US-09-494-190-125	Sequence 125, App
32	119.5	13.5	185	4	US-09-494-190-126	Sequence 126, App
33	118.5	13.4	315	1	US-08-253-155A-34	Sequence 34, Appl
34	118.5	13.4	532	4	US-09-389-956-80	Sequence 80, Appl
35	117	13.2	457	4	US-09-389-956-68	Sequence 68, Appl
36	117	13.2	647	4	US-09-389-956-6	Sequence 6, Appli
37	114.5	13.0	543	4	US-09-362-123A-4	Sequence 4, Appli
38	114.5	13.0	675	1	US-08-317-522A-9	Sequence 9, Appli
39	114.5	13.0	675	1	US-08-439-818A-9	Sequence 9, Appli
40	114.5	13.0	675	2	US-08-751-965-9	Sequence 9, Appli
41	114.5	13.0	675	2	US-08-738-975-9	Sequence 9, Appli
42	114.5	13.0	675	2	US-08-728-626-9	Sequence 9, Appli
43	114.5	13.0	675	3	US-08-808-599A-9	Sequence 9, Appli
44	114	12.9	21	3	US-08-907-468-11	Sequence 11, Appl
45	113.5	12.8	462	3	US-08-486-099-117	Sequence 117, App

ALIGNMENTS

RESULT 1

US-09-087-232A-13

; Sequence 13, Application US/09087232A

; Patent No. 6153431

; GENERAL INFORMATION:

; APPLICANT: Quillent et al.

; TITLE OF INVENTION: HUMAN IMMUNODEFICIENCY VIRUS CO-RECEPTOR

; TITLE OF INVENTION: VARIANTS ASSOCIATED WITH RESISTANCE TO VIRUS INFECTION.

; NUMBER OF SEQUENCES: 23

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: Baker & Botts, L.L.P. attn. Lisa Kole

; STREET: 30 Rockefeller Plaza

; CITY: New York

; STATE: New York

; COUNTRY: USA

; ZIP: 10112

; COMPUTER READABLE FORM:

```

; MEDIUM TYPE: Floppy disk
; COMPUTER: IBM PC compatible
; OPERATING SYSTEM: PC-DOS/MS-DOS
; SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)
; CURRENT APPLICATION DATA:
; APPLICATION NUMBER: US/09/087,232A
; FILING DATE: 28 MAY 1998
; CLASSIFICATION: 435
; PRIOR APPLICATION DATA:
; APPLICATION NUMBER: 60/048,057
; FILING DATE: 30 MAY 1997
; ATTORNEY/AGENT INFORMATION:
; NAME: KOLE, LISA B.
; REGISTRATION NUMBER: 35,225
; REFERENCE/DOCKET NUMBER: AP 31115
; TELECOMMUNICATION INFORMATION:
; TELEPHONE: (212) 408-2628
; TELEFAX: (212) 765-2519
; INFORMATION FOR SEQ ID NO: 13:
; SEQUENCE CHARACTERISTICS:
; LENGTH: 352 amino acids
; TYPE: amino acid
; TOPOLOGY: linear
; MOLECULE TYPE: protein
US-09-087-232A-13

```

```

Query Match          42.9%; Score 379; DB 3; Length 352;
Best Local Similarity 35.0%; Pred. No. 8.1e-35;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db       1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |   |               |||||   | : | : : |
Db      61 LKSMTDIYLLNLAISDLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : ||||||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||||||||||||||           :: |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
          |   |   ||||||||||||||| :|
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 2

```

US-08-861-105-14
; Sequence 14, Application US/08861105
; Patent No. 6258527
; GENERAL INFORMATION:
; APPLICANT: LITTMAN, DAN R.

```

```

; APPLICANT: DENG, HONGKUI
; APPLICANT: ELLMEIER, WILFRIED
; APPLICANT: LANDAU, NATHANIEL R.
; APPLICANT: LIU, RONG
; TITLE OF INVENTION: G-COUPLED RECEPTORS ASSOCIATED WITH
; TITLE OF INVENTION: MACROPHAGE-TROPHIC HIV, AND DIAGNOSTIC AND THERAPEUTIC
; TITLE OF INVENTION: USES THEREOF
; NUMBER OF SEQUENCES: 14
; CORRESPONDENCE ADDRESS:
; ADDRESSEE: David A. Jackson, Esq.
; STREET: 411 Hackensack Ave, Continental Plaza, 4th
; STREET: Floor
; CITY: Hackensack
; STATE: New Jersey
; COUNTRY: USA
; ZIP: 07601
; COMPUTER READABLE FORM:
; MEDIUM TYPE: Floppy disk
; COMPUTER: IBM PC compatible
; OPERATING SYSTEM: PC-DOS/MS-DOS
; SOFTWARE: PatentIn Release #1.0, Version #1.30
; CURRENT APPLICATION DATA:
; APPLICATION NUMBER: US/08/861,105
; FILING DATE:
; CLASSIFICATION: 436
; PRIOR APPLICATION DATA:
; APPLICATION NUMBER: US 08/666,020
; FILING DATE: 19-JUN-1996
; CLASSIFICATION: 436
; PRIOR APPLICATION DATA:
; APPLICATION NUMBER: US 08/227,319
; FILING DATE: 13-APR-1994
; ATTORNEY/AGENT INFORMATION:
; NAME: Jackson Esq., David A.
; REGISTRATION NUMBER: 26,742
; REFERENCE/DOCKET NUMBER: 1049-1-004 N1
; TELECOMMUNICATION INFORMATION:
; TELEPHONE: 201-487-5800
; TELEFAX: 201-343-1684
; INFORMATION FOR SEQ ID NO: 14:
; SEQUENCE CHARACTERISTICS:
; LENGTH: 352 amino acids
; TYPE: amino acid
; STRANDEDNESS: single
; TOPOLOGY: linear
; MOLECULE TYPE: protein
; HYPOTHETICAL: NO
; ORIGINAL SOURCE:
; ORGANISM: Homo sapiens
US-08-861-105-14

```

```

Query Match          42.9%;  Score 379;  DB 3;  Length 352;
Best Local Similarity 35.0%;  Pred. No. 8.1e-35;
Matches 99;  Conservative 7;  Mismatches 27;  Indels 150;  Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
|||||

```


; NAME/KEY: misc_feature
; OTHER INFORMATION: /= "88C amino acid sequence"
US-08-575-967A-2

Query Match 42.9%; Score 379; DB 3; Length 352;
Best Local Similarity 35.0%; Pred. No. 8.1e-35;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          | | |||||||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||||||
Db     119 IILLTIDRYLAVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLLRCRNEKKRHRARLIF 238

Qy     125 -----GGSYKCGLC----QEFGNLCSSSNRLDGHQRV 154
          | | |||||||||||| : |
Db     239 TIMIVYFLFWAPYNIVLLNLTQEFGLNLCSSSNRLDQAMQV 281
```

RESULT 4

US-08-833-752-5

; Sequence 5, Application US/08833752

; Patent No. 6448375

; GENERAL INFORMATION:

; APPLICANT: SAMSON, MICHEL

; APPLICANT: PARMENTIER, MARC

; APPLICANT: VASSART, GILBERT

; APPLICANT: LIBERT, FREDERICK

; TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR

; TITLE OF INVENTION: AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR

; NUMBER OF SEQUENCES: 17

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: Knobbe, Martens, Olson & Bear

; STREET: 620 Newport Center Drive 16th Floor

; CITY: Newport Beach

; STATE: CA

; COUNTRY: U.S.A.

; ZIP: 92660

; COMPUTER READABLE FORM:

; MEDIUM TYPE: Floppy disk

; COMPUTER: IBM PC compatible

; OPERATING SYSTEM: PC-DOS/MS-DOS

; SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

; CURRENT APPLICATION DATA:

; APPLICATION NUMBER: US/08/833,752

; FILING DATE: 9-APR-1997

; CLASSIFICATION: 536


```
; ATTORNEY/AGENT INFORMATION:
; NAME: Altman, Daniel E
; REGISTRATION NUMBER: 34,115
; REFERENCE/DOCKET NUMBER:
; INFORMATION FOR SEQ ID NO: 5:
; SEQUENCE CHARACTERISTICS:
; LENGTH: 352 amino acids
; TYPE: amino acid
; TOPOLOGY: linear
; MOLECULE TYPE: protein
US-08-833-752-5
```

```
Query Match          42.9%; Score 379; DB 4; Length 352;
Best Local Similarity 35.0%; Pred. No. 8.1e-35;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;
```

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      ||||||||||||||||||||||||||||
Db       1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
      | | ||||||||| | : | : : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : |||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      ||||||||||||||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC----QEFG LNNC SSSNRLDGHQRV 154
      | | ||||||||| :|
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFG LNNC SSSNRLDQAMQV 281
```

RESULT 5

US-09-502-783A-2

; Sequence 2, Application US/09502783A

; Patent No. 6511826

; GENERAL INFORMATION:

; APPLICANT: Li, Yi

; APPLICANT: Ruben, Steven M.

; TITLE OF INVENTION: Polynucleotides Encoding Human G-Protein Chemokine Receptor (CCR5)

; TITLE OF INVENTION: HDG NR10

; FILE REFERENCE: 1488.1150006

; CURRENT APPLICATION NUMBER: US/09/502,783A

; CURRENT FILING DATE: 2001-08-23

; PRIOR APPLICATION NUMBER: 08/466,343

; PRIOR FILING DATE: 1995-06-06

; NUMBER OF SEQ ID NOS: 9

; SOFTWARE: PatentIn version 3.0

; SEQ ID NO 2

; LENGTH: 352

; TYPE: PRT

; ORGANISM: Homo sapiens
US-09-502-783A-2

Query Match 42.9%; Score 379; DB 4; Length 352;
Best Local Similarity 35.0%; Pred. No. 8.1e-35;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               ||||||||| | : | : : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||||||||||||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC-----QEFFGLNCCSSSNRLDGHQRV 154
          |   |   ||||||||||||| :|
Db     239 TIMIVYFLFWAPYNIVLLNLTQEFFGLNCCSSSNRLDQAMQV 281
```

RESULT 6

US-09-796-202-1

; Sequence 1, Application US/09796202
; Patent No. 6548636
; GENERAL INFORMATION:
; APPLICANT: Dragic, Tatjana
; APPLICANT: Olson, William
; TITLE OF INVENTION: SULFATED CCR5 PEPTIDES FOR HIV-1 INFECTION
; FILE REFERENCE: 2048/61010/JPW/SHS
; CURRENT APPLICATION NUMBER: US/09/796,202
; CURRENT FILING DATE: 2001-02-28
; NUMBER OF SEQ ID NOS: 17
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 1
; LENGTH: 352
; TYPE: PRT
; ORGANISM: human
US-09-796-202-1

Query Match 42.9%; Score 379; DB 4; Length 352;
Best Local Similarity 35.0%; Pred. No. 8.1e-35;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               ||||||||| | : | : : |
```

Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||||
 Db 119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 : |||||
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCLC----QEFG LNNC SSSNRLDGHQRV 154
 : |||||
 Db 239 TIMIVYFLFWAPYNIVLLNTFQEFG LNNC SSSNRLDQAMQV 281

RESULT 7

US-09-045-583-52

; Sequence 52, Application US/09045583

; Patent No. 6287805

; GENERAL INFORMATION:

; APPLICANT: Graham, Gerard J. et al.

; TITLE OF INVENTION: No. 6287805el Molecules of the G Protein-Coupled

; NUMBER OF SEQUENCES: 56

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: LAHIVE & COCKFIELD, LLP

; STREET: 28 State Street

; CITY: Boston

; STATE: Massachusetts

; COUNTRY: USA

; ZIP: 02109

; COMPUTER READABLE FORM:

; MEDIUM TYPE: Floppy disk

; COMPUTER: IBM PC compatible

; OPERATING SYSTEM: PC-DOS/MS-DOS

; SOFTWARE: PatentIn Release #1.0, Version #1.25

; CURRENT APPLICATION DATA:

; APPLICATION NUMBER: US/09/045,583

; FILING DATE: 20-MAR-98

; CLASSIFICATION: 435

; PRIOR APPLICATION DATA:

; APPLICATION NUMBER:

; FILING DATE:

; ATTORNEY/AGENT INFORMATION:

; NAME: Mandragouras, Amy E.

; REGISTRATION NUMBER: 36,207

; REFERENCE/DOCKET NUMBER: MNI-044

; TELECOMMUNICATION INFORMATION:

; TELEPHONE: (617)227-7400

; TELEFAX: (617)742-4214

; INFORMATION FOR SEQ ID NO: 52:

; SEQUENCE CHARACTERISTICS:

; LENGTH: 352 amino acids

; TYPE: amino acid

; TOPOLOGY: linear

; MOLECULE TYPE: peptide

; FRAGMENT TYPE: internal

US-09-045-583-52

Query Match 42.3%; Score 374; DB 3; Length 352;
Best Local Similarity 34.6%; Pred. No. 3e-34;
Matches 98; Conservative 8; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      |||||||||:|||||||||||
Db      1 MDYQVSSPIYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | ||||||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : |||||||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      ||||||||| : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
      | | ||||||||| : |
Db      239 TIMIVYFLFWAPYNIVLLNTFOEFFGLNNCSSSNRLDOAMOV 281

```

RESULT 8

US-09-534-185-52

; Sequence 52, Application US/09534185

; Patent No. 6403767

; GENERAL INFORMATION:

; APPLICANT: Graham, Gerard J. et al.

; TITLE OF INVENTION: No. 6403767e1 Molecules of the G Protein-Coupled
 ; Heptahelical Receptor Superfamily and Uses
 ; Therefor

; NUMBER OF SEQUENCES: 56

CORRESPONDENCE ADDRESS:

ADDRESSEE: LAHIVE & COCKFIELD, LLP

STREET: 28 State Street

; CITY: Boston

; STATE: Massachusetts

; COUNTRY: USA

; ZIP: 02109

; COMPUTER READABLE FORM:

```
; MEDIUM TYPE: Floppy disk
```

```

;          COMPUTER: IBM PC compatible

```

OPERATING SYSTEM: PC-DOS/MS-DOS

```
; SOFTWARE: PatentIn Release #1.0, Version #1.25
```

```

; CURRENT APPLICATION DATA:

```

APPLICATION NUMBER: US/09/534,185

FILING DATE: 24-Mar-2000

; CLASSIFICATION: <Unknown>

; PRIOR APPLICATION DATA:

; APPLICATION NUMBER: 09/045,583

; FILING DATE: <Unknown>

; ATTORNEY/AGENT INFORMATION:

NAME: Mandragouras, Amy E.

```

;          REGISTRATION NUMBER: 36,207
;          REFERENCE/DOCKET NUMBER: MNI-044
;          TELECOMMUNICATION INFORMATION:
;          TELEPHONE: (617)227-7400
;          TELEFAX: (617)742-4214
;          INFORMATION FOR SEQ ID NO: 52:
;          SEQUENCE CHARACTERISTICS:
;          LENGTH: 352 amino acids
;          TYPE: amino acid
;          TOPOLOGY: linear
;          MOLECULE TYPE: peptide
;          FRAGMENT TYPE: internal
;          SEQUENCE DESCRIPTION: SEQ ID NO: 52:
US-09-534-185-52

```

```

Query Match          42.3%;  Score 374;  DB 4;  Length 352;
Best Local Similarity 34.6%;  Pred. No. 3e-34;
Matches   98;  Conservative   8;  Mismatches   27;  Indels   150;  Gaps    7;

```

```

Qy      20 MDYQVSSPIYDINYTSEPCQKINVKQIAA----- 49
          |||||:|||||
Db      1 MDYQVSSPIYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | :  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||                      :: |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLLRCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  |                      |||||  :|
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 9

US-08-466-343D-2

; Sequence 2, Application US/08466343D

; Patent No. 6025154

; GENERAL INFORMATION:

; APPLICANT: LI, Yi

; TITLE OF INVENTION: POLYNUCLEOTIDES ENCODING HUMAN G-PROTEIN

; TITLE OF INVENTION: CHEMOKINE RECEPTOR HDG NR10 (AS AMENDED)

; NUMBER OF SEQUENCES: 9

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

; STREET: 1100 NEW YORK AVE., NW, SUITE 600

; CITY: WASHINGTON

; STATE: DC

; COUNTRY: USA

; ZIP: 20005

; COMPUTER READABLE FORM:

```

; MEDIUM TYPE: Floppy disk
; COMPUTER: IBM PC compatible
; OPERATING SYSTEM: PC-DOS/MS-DOS
; SOFTWARE: PatentIn Release #1.0, Version #1.30
; CURRENT APPLICATION DATA:
; APPLICATION NUMBER: US/08/466,343D
; FILING DATE: 06-JUN-1995
; CLASSIFICATION: 435
; ATTORNEY/AGENT INFORMATION:
; NAME: STEFFE, ERIC K.
; REGISTRATION NUMBER: 36,688
; REFERENCE/DOCKET NUMBER: 1488.1150000/EKS/KLM
; TELECOMMUNICATION INFORMATION:
; TELEPHONE: (202) 371-2600
; TELEFAX: (202) 371-2540
; INFORMATION FOR SEQ ID NO: 2:
; SEQUENCE CHARACTERISTICS:
; LENGTH: 352 amino acids
; TYPE: amino acid
; TOPOLOGY: linear
; MOLECULE TYPE: protein
US-08-466-343D-2

```

```

Query Match          42.2%; Score 373; DB 3; Length 352;
Best Local Similarity 34.6%; Pred. No. 3.9e-34;
Matches 98; Conservative 7; Mismatches 28; Indels 150; Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||||||||||||| |||||
Db      1 MDYQVSSPIYDINYYTSEPCPKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCQR 60

Qy      50 -----YKGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          | | ||||||||| | : | : |
Db      61 LESMTDIYLLNLAISDLFFLLTVPFWAHAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||| : : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKGLC----QEFFGLNNCSSSNRLDGHQRV 154
          | | ||||||||| :|
Db      239 TIMIVYFLFWAPYNI VLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 10

US-09-517-605-5

```

; Sequence 5, Application US/09517605
; Patent No. 6391567
; GENERAL INFORMATION:
; APPLICANT: Littman, Dan R.
; APPLICANT: Kwon, Douglas S.
; APPLICANT: van Kooyk, Yvette
; APPLICANT: Geijtenbeck, Theo

```

```
; TITLE OF INVENTION: METHODS OF USING A FACILITATOR OF RETROVIRAL ENTRY INTO
; TITLE OF INVENTION: CELLS
; FILE REFERENCE: 1049-1-017
; CURRENT APPLICATION NUMBER: US/09/517,605
; CURRENT FILING DATE: 2000-03-02
; NUMBER OF SEQ ID NOS: 17
; SOFTWARE: PatentIn Ver. 2.0
; SEQ ID NO 5
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Homo sapiens
US-09-517-605-5
```

```
Query Match          40.6%; Score 359; DB 4; Length 352;
Best Local Similarity 33.9%; Pred. No. 1.5e-32;
Matches 96; Conservative 8; Mismatches 29; Indels 150; Gaps 7;
```

```
Qy      20 MDYQVSSPIYDINYTTSEPCQKINVKQIAA----- 49
          ||||| |||:| |||||
Db      1 MDYQVSSPTYDIDYDTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILVLINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |  | ||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||| ||||| : : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  | ||||| :|
Db      239 TIMIVYFLFWAPYNIVLLNLTQEFFGLNNCSSSNRLDQAMQV 281
```

RESULT 11

US-08-724-984A-2

; Sequence 2, Application US/08724984A

; Patent No. 6388055

; GENERAL INFORMATION:

; APPLICANT: Derk Bergsma, Mary Brawner, and Usman Shabon

; TITLE OF INVENTION: No. 6388055el Mouse Genomic Clone of the CC-

; TITLE OF INVENTION: CKR5 Receptor

; NUMBER OF SEQUENCES: 5

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: SmithKline Beecham Corporation

; STREET: 709 Swedeland Road, P.O. Box 1539

; CITY: King of Prussia

; STATE: PA

; COUNTRY: USA

; ZIP: 19406-0939

; COMPUTER READABLE FORM:

; MEDIUM TYPE: DISKETTE, 3.5 INCH, 1.44 Mb STORAGE

; COMPUTER: IBM 486

```

;   OPERATING SYSTEM:  WINDOWS FOR WORKGROUPS
;   SOFTWARE:  MICROSOFT WORD
;   CURRENT APPLICATION DATA:
;   APPLICATION NUMBER:  US/08/724,984A
;   FILING DATE:  October 3, 1996
;   CLASSIFICATION:  800
;   PRIOR APPLICATION DATA:
;   APPLICATION NUMBER:
;   FILING DATE:
;   ATTORNEY/AGENT INFORMATION:
;   NAME:  William T. Han
;   REGISTRATION NUMBER:  34,344
;   REFERENCE/DOCKET NUMBER:  ATG50023
;   TELECOMMUNICATION INFORMATION:
;   TELEPHONE:  610 270 5024
;   TELEFAX:  610 270 5090
;   INFORMATION FOR SEQ ID NO:  2:
;   SEQUENCE CHARACTERISTICS:
;   LENGTH:  354
;   TYPE:  Amino Acid
;   TOPOLOGY:  Linear
US-08-724-984A-2

```

```

Query Match          30.8%;  Score 272;  DB 4;  Length 354;
Best Local Similarity 27.9%;  Pred. No. 1e-22;
Matches   80;  Conservative   12;  Mismatches   29;  Indels  166;  Gaps    8;

```

```

Qy      20 MDYQVSSP--IYDINYTSEPCQKINVKQIAA----- 49
      ||:| | | | ||||:| | |||||
Db      1 MDFQGSVPTYIYDIDYGMSAPCQKINVKQIAAQLLPPLYSLVFIFGFAGNMMVFLILISC 60

Qy      50 -----YKGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG- 83
      | | | | | | | | | | | | | | | | | | | | | |
Db      61 KKLKSVTDIYLLNLAISDLLFLLTLPFWAHYAANEWIFGNIMC--KVFTGVYHIGYFGGI 118

Qy      84 -----LCTRSQKEGLHY 95
      : ||||| | |
Db      119 FFIILLTIDRYLAIVHAVFALKVTTVNFGVITSVVTWVAVFASLPEIIFTRSQKEGFHY 178

Qy      96 TCSSHFPYSQYQFWKNFQTLK-----IHQRVHGGGGSYKC----- 130
      ||| |||:| | |||:|||| | | | | | | | | | | | | | | | | | | | | | |
Db      179 TCSPHFPHTQYHFWKSFQTLKMVILSLILPLLMIICYSGLIHT-----LFRCRNEKK 231

Qy      131 -----GLCQEFFGLNNCSSSNRLD 149
      |||||
Db      232 RHRAVRLIFAIMIVYFLFWTPYNIVLLLTTFQEFFGLNNCSSSNRLD 278

```

RESULT 12

US-08-833-752-4

; Sequence 4, Application US/08833752

; Patent No. 6448375

; GENERAL INFORMATION:

```

;   APPLICANT:  SAMSON, MICHEL
;   APPLICANT:  PARMENTIER, MARC
;   APPLICANT:  VASSART, GILBERT
;   APPLICANT:  LIBERT, FREDERICK

```



```

; TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR
; TITLE OF INVENTION: AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR
; NUMBER OF SEQUENCES: 17
; CORRESPONDENCE ADDRESS:
; ADDRESSEE: Knobbe, Martens, Olson & Bear
; STREET: 620 Newport Center Drive 16th Floor
; CITY: Newport Beach
; STATE: CA
; COUNTRY: U.S.A.
; ZIP: 92660
; COMPUTER READABLE FORM:
; MEDIUM TYPE: Floppy disk
; COMPUTER: IBM PC compatible
; OPERATING SYSTEM: PC-DOS/MS-DOS
; SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)
; CURRENT APPLICATION DATA:
; APPLICATION NUMBER: US/08/833,752
; FILING DATE: 9-APR-1997
; CLASSIFICATION: 536
; ATTORNEY/AGENT INFORMATION:
; NAME: Altman, Daniel E
; REGISTRATION NUMBER: 34,115
; REFERENCE/DOCKET NUMBER:
; INFORMATION FOR SEQ ID NO: 4:
; SEQUENCE CHARACTERISTICS:
; LENGTH: 184 amino acids
; TYPE: amino acid
; TOPOLOGY: linear
; MOLECULE TYPE: protein
US-08-833-752-4

```

```

Query Match          29.2%; Score 258; DB 4; Length 184;
Best Local Similarity 34.9%; Pred. No. 1.8e-21;
Matches 65; Conservative 4; Mismatches 13; Indels 104; Gaps 4;

```

```

Qy      20 MDYQVSSPIYDINYTTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYTTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPY 103
          |||||
Db      179 SSHFPY 184

```

```

RESULT 13
US-09-087-232A-17
; Sequence 17, Application US/09087232A
; Patent No. 6153431
; GENERAL INFORMATION:

```

```

;   APPLICANT:  Quillent et al.
;   TITLE OF INVENTION:  HUMAN IMMUNODEFICIENCY VIRUS CO-RECEPTOR
;   TITLE OF INVENTION:  VARIANTS ASSOCIATED WITH RESISTANCE TO VIRUS
INFECTION.
;   NUMBER OF SEQUENCES:  23
;   CORRESPONDENCE ADDRESS:
;       ADDRESSEE:  Baker & Botts, L.L.P. attn. Lisa Kole
;       STREET:  30 Rockefeller Plaza
;       CITY:  New York
;       STATE:  New York
;       COUNTRY:  USA
;       ZIP:  10112
;   COMPUTER READABLE FORM:
;       MEDIUM TYPE:  Floppy disk
;       COMPUTER:  IBM PC compatible
;       OPERATING SYSTEM:  PC-DOS/MS-DOS
;       SOFTWARE:  PatentIn Release #1.0, Version #1.30 (EPO)
;   CURRENT APPLICATION DATA:
;       APPLICATION NUMBER:  US/09/087,232A
;       FILING DATE:  28 MAY 1998
;       CLASSIFICATION:  435
;   PRIOR APPLICATION DATA:
;       APPLICATION NUMBER:  60/048,057
;       FILING DATE:  30 MAY 1997
;   ATTORNEY/AGENT INFORMATION:
;       NAME:  KOLE, LISA B.
;       REGISTRATION NUMBER:  35,225
;       REFERENCE/DOCKET NUMBER:  AP 31115
;   TELECOMMUNICATION INFORMATION:
;       TELEPHONE:  (212) 408-2628
;       TELEFAX:  (212) 765-2519
;   INFORMATION FOR SEQ ID NO:  17:
;       SEQUENCE CHARACTERISTICS:
;           LENGTH: 215 amino acids
;           TYPE:  amino acid
;           TOPOLOGY:  linear
;       MOLECULE TYPE:  protein
US-09-087-232A-17

```

```

Query Match          29.2%;  Score 258;  DB 3;  Length 215;
Best Local Similarity 34.9%;  Pred. No. 2.2e-21;
Matches 65;  Conservative 4;  Mismatches 13;  Indels 104;  Gaps 4;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | :  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPY 103
          |||||

```

Db 179 SSHFPY 184

RESULT 14

US-08-833-752-6

; Sequence 6, Application US/08833752

; Patent No. 6448375

; GENERAL INFORMATION:

; APPLICANT: SAMSON, MICHEL

; APPLICANT: PARMENTIER, MARC

; APPLICANT: VASSART, GILBERT

; APPLICANT: LIBERT, FREDERICK

; TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR

; TITLE OF INVENTION: AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR

; NUMBER OF SEQUENCES: 17

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: Knobbe, Martens, Olson & Bear

; STREET: 620 Newport Center Drive 16th Floor

; CITY: Newport Beach

; STATE: CA

; COUNTRY: U.S.A.

; ZIP: 92660

; COMPUTER READABLE FORM:

; MEDIUM TYPE: Floppy disk

; COMPUTER: IBM PC compatible

; OPERATING SYSTEM: PC-DOS/MS-DOS

; SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

; CURRENT APPLICATION DATA:

; APPLICATION NUMBER: US/08/833,752

; FILING DATE: 9-APR-1997

; CLASSIFICATION: 536

; ATTORNEY/AGENT INFORMATION:

; NAME: Altman, Daniel E

; REGISTRATION NUMBER: 34,115

; REFERENCE/DOCKET NUMBER:

; INFORMATION FOR SEQ ID NO: 6:

; SEQUENCE CHARACTERISTICS:

; LENGTH: 215 amino acids

; TYPE: amino acid

; TOPOLOGY: linear

; MOLECULE TYPE: protein

US-08-833-752-6

Query Match 29.2%; Score 258; DB 4; Length 215;

Best Local Similarity 34.9%; Pred. No. 2.2e-21;

Matches 65; Conservative 4; Mismatches 13; Indels 104; Gaps 4;

```
QY      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db       1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

QY      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          | |                      ||||| | : | : : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

QY      84 -----LCTRSQKEGLHYTC 97
          : |||||||||||
```


Db 1 MDYQVSSPIYDINYTTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
Qy 50 -----YKCGLC-----AAAQWDFGNTM 66
 | | |||||||
Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTM 100

Search completed: March 5, 2004, 16:30:39
Job time : 43.1574 secs

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:16:19 ; Search time 35.858 Seconds
(without alignments)
421.163 Million cell updates/sec

Title: US-10-057-890A-31
Perfect score: 884
Sequence: 1 MKVSVAALSCIMLV TALGSM.....GLNNCSSSNRLDGHQRVHAA 157

Scoring table: BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 283366 seqs, 96191526 residues

Total number of hits satisfying chosen parameters: 283366

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : PIR_78:*
1: pir1:*
2: pir2:*
3: pir3:*
4: pir4:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Query		DB	ID	Description
	Score	Match Length			
1	379	42.9	352	2	A43113 chemokine (C-C) re
2	154	17.4	201	2	I57505 zinc finger protei
3	144	16.3	693	2	I37570 zinc finger protei
4	143.5	16.2	319	2	T46469 hypothetical prote
5	141	16.0	555	2	I53869 zinc finger protei
6	138.5	15.7	223	2	F31201 GLI-related finger
7	136.5	15.4	348	2	I38599 zinc finger protei
8	136.5	15.4	386	2	T12527 hypothetical prote
9	135.5	15.3	572	2	I39311 Kruppel-type zinc
10	134	15.2	710	2	I48668 zinc finger protei
11	133.5	15.1	475	2	S03679 finger protein (cl
12	133.5	15.1	636	2	I48689 gene NK10 protein
13	133.5	15.1	701	2	T14757 hypothetical prote

14	131.5	14.9	686	2	A34612	zinc finger protei
15	131	14.8	261	2	S70006	finger protein zfo
16	129.5	14.6	614	2	JH0500	zinc finger protei
17	128.5	14.5	580	2	A37107	spermatogenesis pr
18	127.5	14.4	469	2	I38600	zinc finger protei
19	127.5	14.4	1173	2	I50620	procKr2 - chicken
20	127	14.4	728	2	A48830	probable transcrip
21	126.5	14.3	732	2	S47073	finger protein HZF
22	124	14.0	367	2	S06582	finger protein (cl
23	124	14.0	589	2	I38598	zinc finger protei
24	123.5	14.0	209	2	S47068	finger protein HZF
25	123.5	14.0	347	2	S00549	developmental cont
26	123.5	14.0	378	2	S33994	finger protein ZNF
27	122.5	13.9	488	2	S47072	finger protein HZF
28	122.5	13.9	594	2	T12488	hypothetical prote
29	122.5	13.9	803	2	S26823	zinc finger protei
30	122	13.8	1350	2	S00647	finger protein - A
31	121.5	13.7	337	2	S60520	finger protein ZNF
32	120.5	13.6	169	2	A39240	finger protein mfg
33	120.5	13.6	229	2	A48927	Kruppel-like zinc
34	120.5	13.6	292	2	S43826	finger protein OZF
35	120.5	13.6	399	2	S47071	finger protein HZF
36	120.5	13.6	546	2	I49636	DNA-binding protei
37	120.5	13.6	1191	2	S35305	zinc finger protei
38	120	13.6	435	2	S00833	finger protein (cl
39	120	13.6	654	2	A57785	finger protein ZNF
40	119.5	13.5	194	2	I53859	zinc finger protei
41	119.5	13.5	393	2	JN0533	finger protein pML
42	119.5	13.5	576	2	A48157	renal transcriptio
43	119.5	13.5	671	2	JE0288	krueppel-type zinc
44	118.5	13.4	107	2	I39315	zinc-finger protei
45	118.5	13.4	120	2	G02493	KR-ZNF1 - human (f

ALIGNMENTS

RESULT 1

A43113

chemokine (C-C) receptor 5 - human

N;Alternate names: C-C CKR-5; CCR5

C;Species: Homo sapiens (man)

C;Date: 12-Jul-1996 #sequence_revision 12-Jul-1996 #text_change 20-Jun-2000

C;Accession: A43113; S71808; A58834; A58832; G02653; A58833

R;Samson, M.; Labbe, O.; Mollereau, C.; Vassart, G.; Parmentier, M.

Biochemistry 35, 3362-3367, 1996

A;Title: Molecular cloning and functional expression of a new human CC-chemokine receptor gene.

A;Reference number: A43113; MUID:96241590; PMID:8639485

A;Accession: A43113

A;Molecule type: mRNA

A;Residues: 1-352 <SAM1>

A;Cross-references: GB:X91492; NID:g1262810; PIDN:CAA62796.1; PID:g1262811

R;Samson, M.; Libert, F.; Doranz, B.J.; Rucker, J.; Liesnard, C.; Farber, C.M.; Saragosti, S.; Lapoumeroulie, C.; Cognaux, J.; Forceille, C.; Muyldermans, G.; Verhofstede, C.; Burtonboy, G.; Georges, M.; Imai, T.; Rana, S.; Yi, Y.; Smyth, R.J.; Collman, R.G.; Doms, R.W.; Vassart, G.; Parmentier, M.

Nature 382, 722-725, 1996

A;Title: Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene.

A;Reference number: S71808; MUID:96345670; PMID:8751444

A;Accession: S71808

A;Status: nucleic acid sequence not shown; not compared with conceptual translation

A;Molecule type: DNA

A;Residues: 182-206;207-230 <SAM2>

A;Accession: A58834

A;Status: nucleic acid sequence not shown; not compared with conceptual translation

A;Molecule type: DNA

A;Residues: 1-184,'IKDShLGAGPAAACHGHLLLLGNPKNSASVSK' <SAM3>

A;Cross-references: GB:X99393; NID:g1524062; PIDN:CAA67767.1; PID:g1524063

A;Note: this frameshift mutation results in a non-functional receptor but confers a degree of resistance to HIV-1 infection; it has an allele frequency of 0.09 or more in some caucasian populations and may have had a selective advantage by conferring resistance to Yersinia plague infections

R;Combadiere, C.; Ahuja, S.K.; Tiffany, H.L.; Murphy, P.M.

J. Leukoc. Biol. 60, 147-152, 1996

A;Title: Cloning and functional expression of CC CKR5, a human monocyte CC chemokine receptor selective for MIP-1alpha, MIP-1beta, and RANTES.

A;Reference number: A58832; MUID:96295970; PMID:8699119

A;Accession: A58832

A;Molecule type: mRNA

A;Residues: 1-352 <COM1>

A;Cross-references: GB:U57840; NID:g1502408; PIDN:AAB17071.1; PID:g1502409

A;Experimental source: clone 8, endotoxin-stimulated peripheral blood monocytes

R;Combadiere, C.

submitted to the EMBL Data Library, May 1996

A;Reference number: H01541

A;Accession: G02653

A;Status: translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-89,'L',91-352 <COM2>

A;Cross-references: EMBL:U57840

R;Raport, C.J.; Gosling, J.; Schweickart, V.L.; Gray, P.W.; Charo, I.F.

J. Biol. Chem. 271, 17161-17166, 1996

A;Title: Molecular cloning and functional characterization of a novel human CC chemokine receptor (CCR5) for RANTES, MIP-1beta, and MIP-1alpha.

A;Reference number: A58833; MUID:96291862; PMID:8663314

A;Accession: A58833

A;Molecule type: mRNA

A;Residues: 1-352 <RAP>

A;Cross-references: GB:U54994; NID:g1457945; PIDN:AAC50598.1; PID:g1457946

C;Comment: This is a receptor for chemokines MIP-1alpha (see PIR:A30574), MIP-1beta (see PIR:A31767), and RANTES (see PIR:A28815).

C;Comment: Macrophage- and dual-tropic strains of HIV-1 bind to a complex of chemokine (C-C) receptor 5 and T-cell surface glycoprotein CD4 (see PIR:RWHUT4).

C;Genetics:

A;Gene: GDB:CMKBR5; CCR5; CKR-5; CC-CKR-5; CKR5; ChemR13

A;Cross-references: GDB:1230510; OMIM:601373

A;Map position: 3p21-3p21

C;Function:

A;Description: G protein-coupled receptor for chemokines MIP-1alpha, MIP-1beta and RANTES

A;Note: probably acts to control granulocyte proliferation and differentiation
 C;Superfamily: vertebrate rhodopsin
 C;Keywords: AIDS; G protein-coupled receptor; glycoprotein; phosphoprotein; transmembrane protein
 F;32-56/Domain: transmembrane #status predicted <TM1>
 F;67-87/Domain: transmembrane #status predicted <TM2>
 F;103-124/Domain: transmembrane #status predicted <TM3>
 F;142-166/Domain: transmembrane #status predicted <TM4>
 F;193-218/Domain: transmembrane #status predicted <TM5>
 F;236-257/Domain: transmembrane #status predicted <TM6>
 F;285-300/Domain: transmembrane #status predicted <TM7>
 F;20-269,101-178/Disulfide bonds: #status predicted
 F;268/Binding site: carbohydrate (Asn) (covalent) #status predicted
 F;336,337,342/Binding site: phosphate (Ser) (covalent) #status predicted
 F;340,343/Binding site: phosphate (Thr) (covalent) #status predicted

Query Match 42.9%; Score 379; DB 2; Length 352;
 Best Local Similarity 35.0%; Pred. No. 8e-29;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               |||
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYS GILKTL LRCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154
          |   |               |||
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281
  
```

RESULT 2

I57505

zinc finger protein - mouse

C;Species: Mus musculus (house mouse)

C;Date: 02-Aug-1996 #sequence_revision 02-Aug-1996 #text_change 05-Nov-1999

C;Accession: I57505

R;Ernault-Lange, M.; Kress, M.; Hamer, D.

Mol. Cell. Biol. 10, 418-421, 1990

A;Title: A gene that encodes a protein consisting solely of zinc finger domains is preferentially expressed in transformed mouse cells.

A;Reference number: I57505; MUID:90097859; PMID:2104662

A;Accession: I57505

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-201 <RES>

A;Cross-references: GB:M32057; NID:g199766; PIDN:AAA39729.1; PID:g199767

C;Genetics:
A;Gene: MOK2

Query Match 17.4%; Score 154; DB 2; Length 201;
Best Local Similarity 37.0%; Pred. No. 1.6e-07;
Matches 44; Conservative 13; Mismatches 44; Indels 18; Gaps 6;

```
Qy      45 KQIAAYKCGLCAAAQWDFGNTMCQHQRVHGHHHSYKCGLCTR--SQKEGLHYTCSSHF- 101
      | | :||: : : : : ||||| |||| | : || || :|
Db      60 KPYACEECGMSFSQR----SNLHIHQRVH-TGERPYKCGECGKGFSQSSNLHIHRCTHTG 114

Qy     102 --PYSQYQFWKNF---QTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
      || |: | | :|| ||| | | || | : | | |:| |||||
Db     115 EKPYQCYECGKGFSQSSDLRIHLRVHTGEKPYHCGKCGQGF-----SQSSKLLIHQRVH 168
```

RESULT 3

I37570

zinc finger protein - human (fragment)

C;Species: Homo sapiens (man)

C;Date: 29-May-1998 #sequence_revision 29-May-1998 #text_change 01-Dec-2000

C;Accession: I37570

R;Abrink, M.; Aveskogh, M.; Hellman, L.

DNA Cell Biol. 14, 125-136, 1995

A;Title: Isolation of cDNA clones for 42 different Kruppel-related zinc finger proteins expressed in the human monoblast cell line U-937.

A;Reference number: I37566; MUID:95169271; PMID:7865130

A;Accession: I37570

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-693 <RES>

A;Cross-references: EMBL:X78927; NID:g498726; PIDN:CAA55527.1; PID:g498727

C;Genetics:

A;Gene: HZF4

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 16.3%; Score 144; DB 2; Length 693;
Best Local Similarity 33.3%; Pred. No. 5.1e-06;
Matches 39; Conservative 15; Mismatches 33; Indels 30; Gaps 7;

```
Qy      40 QKINVKQIAAYKCGLCAAA-QWDFGNTMCQHQRVHGHHHSYKCGLCTRSQKEGLHYTCS 98
      ||:: : |||| | :| | |||| | || | | :: :
Db     554 QKVHTGE-KPYKCGECGKGFKWSLNLDLDM--HQRVH-TGEKPYTCGAC-----GKHFSQA 603

Qy      99 SHFPYSQYQFWKNFQTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
      | :::|| || | ||| :| : | | |:| |:|||
Db     604 S-----SLQLHQSVHTGEKPYKCDVCGKVF-----SRSSQLQYHRRVH 641
```

RESULT 4

T46469

hypothetical protein DKFZp434G1930.1 - human (fragment)

C;Species: Homo sapiens (man)

C;Date: 04-Feb-2000 #sequence_revision 04-Feb-2000 #text_change 04-Feb-2000

C;Accession: T46469

R;Blum, H.; Bauersachs, S.; Mewes, H.W.; Gassenhuber, J.; Wiemann, S.

submitted to the Protein Sequence Database, January 2000

A;Reference number: Z23034
 A;Accession: T46469
 A;Status: preliminary
 A;Molecule type: mRNA
 A;Residues: 1-319 <AAA>
 A;Cross-references: EMBL:AL137483
 A;Experimental source: adult testis; clone DKFZp434G1930
 C;Genetics:
 A;Note: DKFZp434G1930.1

Query Match 16.2%; Score 143.5; DB 2; Length 319;
 Best Local Similarity 32.8%; Pred. No. 2.6e-06;
 Matches 38; Conservative 15; Mismatches 44; Indels 19; Gaps 7;

```

Qy      50 YKCGLCAAAQWDFGNTMCQHQRVHGHHSYKCGLCTRSQKEGLHYTCSSHF-----P 102
      ||| | : :: |||:| ||| | : | |
Db      9 YCNECEKT-FSHRSSLLSHQRIH-TGEKPYKCNECEKAFSNS--STLIKHLRVHTGEKP 64

Qy     103 YSQYQFWKNF---QTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSNRLDGHQRVH 155
      | : | | || :|||:| | |||| |:: | | : : | |||:|
Db     65 YRCRECGKAFSQCSTLTVHQRIHTGEKLYKCGECEKAF---NCRA--KLHRHQRIH 115
  
```

RESULT 5

I53869

zinc finger protein - mouse

C;Species: Mus musculus (house mouse)

C;Date: 29-May-1998 #sequence_revision 29-May-1998 #text_change 01-Dec-2000

C;Accession: I53869

R;Brady, J.P.; Piatigorsky, J.

Gene 149, 299-304, 1994

A;Title: A mouse cDNA encoding a protein with zinc-fingers and a KRAB domain shows similarity to human profilaggrin.

A;Reference number: I53869; MUID:95047492; PMID:7959006

A;Accession: I53869

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-555 <RES>

A;Cross-references: GB:L28167; NID:g758660; PIDN:AAA67545.1; PID:g758661

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 16.0%; Score 141; DB 2; Length 555;
 Best Local Similarity 23.8%; Pred. No. 7.9e-06;
 Matches 50; Conservative 25; Mismatches 47; Indels 88; Gaps 11;

```

Qy      23 QVSSPIYDINYYTSEPCQ-----KINVKQIAAYKCGLCAAAQWDFGNTMCQH 69
      |:| :| : ||| |:| |:: :: | | | :| : : : |
Db     303 RVCTP---VKPYTCEQCEKSLLDVQHLMSHVKVHTRE-RPYNCETCGSA-FSQASHLQDH 357

Qy      70 QRVH-----GHHHHS-----YKCGLCTRSQKEGLHYTCSSHF- 101
      ||:| | | | | |||| | :| : |||:
Db     358 QRLHTGEKPFKCDACGKSFSRSSHLRSHQRVHTGEKPYKCGECGKS-----FICSSNLY 411

Qy     102 -----PYSQYQFWKNF---QTLKIHQRVHGGGGSYKCGLCQEFFGLNN----- 141
      || | | | :| :|| :| | || | :| : : ||:
Db     412 IHQRVHTGEKPYKCVDCGKEFSRPSLQAHQGIHTGEKSIVCTMCGKGYTLNSNLQVHLR 471
  
```

Qy 142 -----C-----SSSNRLDGHQRVH 155
 | | | : | | | |
 Db 472 VHTGEKPYSCDVCCKGKFSRSSQLQSHQRVH 501

RESULT 6

F31201

GLI-related finger protein HKR4 - human (fragments)

C;Species: Homo sapiens (man)

C;Date: 31-Mar-1990 #sequence_revision 31-Mar-1990 #text_change 17-Mar-1994

C;Accession: F31201

R;Ruppert, J.M.; Kinzler, K.W.; Wong, A.J.; Bigner, S.H.; Kao, F.T.; Law, M.L.; Seuanez, H.N.; O'Brien, S.J.; Vogelstein, B.

Mol. Cell. Biol. 8, 3104-3113, 1988

A;Title: The GLI-Kruppel family of human genes.

A;Reference number: A93103; MUID:89096896; PMID:2850480

A;Accession: F31201

A;Molecule type: DNA

A;Residues: 1-223 <RUP>

C;Keywords: DNA binding; zinc finger

Query Match 15.7%; Score 138.5; DB 2; Length 223;
 Best Local Similarity 30.7%; Pred. No. 5.6e-06;
 Matches 42; Conservative 13; Mismatches 39; Indels 43; Gaps 7;

Qy 50 YKCGLCAAAQWDFGNTMCQHQRVHGHHSYKCGLCTRSQKEGLHYTCSSHFYPYSQYQFW 109
 | : | | | : : | | | : | | | | | : | | : | | : |

Db 96 YECGQCGRF-FSHSSHFTQHLRIH-NGEKPYKCGECVRHQ--LH---TGEKPYACSQCG 148

Qy 110 KNF---QTLKIHQRVHGGGGSYK-----CGLCQEFG 138
 | | | | | : | | : | | : | | : | |

Db 149 KAFIWSSVLIEHQRIHTGEKPYECSDCGKAFRGRSHFFRHLRTHTGEKPFACGACGKAFG 208

Qy 139 LNNCSSSNRLDGHQRVH 155

| : | | | |
 Db 209 -----QSSQLIQHQRVH 220

RESULT 7

I38599

zinc finger protein ZNF134 - human

C;Species: Homo sapiens (man)

C;Date: 23-Feb-1996 #sequence_revision 23-Feb-1996 #text_change 05-Nov-1999

C;Accession: I38599

R;Tommerup, N.; Vissing, H.

Genomics 27, 259-264, 1995

A;Title: Isolation and fine mapping of 16 novel human zinc finger-encoding cDNAs identify putative candidate genes for developmental and malignant disorders.

A;Reference number: A57785; MUID:96044430; PMID:7557990

A;Accession: I38599

A;Status: preliminary

A;Molecule type: mRNA

A;Residues: 1-348 <RES>

A;Cross-references: EMBL:U09412; NID:g488552; PIDN:AAC50253.1; PID:g488553

C;Genetics:

A;Gene: GDB:ZNF134

A;Cross-references: GDB:137033

A;Map position: 19q13.4-19q13.4

Query Match 15.4%; Score 136.5; DB 2; Length 348;
Best Local Similarity 27.3%; Pred. No. 1.3e-05;
Matches 45; Conservative 13; Mismatches 46; Indels 61; Gaps 9;

```
Qy      50 YKCGLCAAAQWDFGNTMCQHQRVHGHHSYKCGLC-----TRSQKEGLH-----YT 96
      ||| | | : :|: |||:| | :| | | : :| |
Db      97 YKCSECGKA-FSRKDTLVQHQRHS-GEKPYECSECGKAFSRKATLVQHQRHTGERPYE 154

Qy      97 CS-----SHFPYSQYQFWKNF---QTLKIHQRVHGGGGSYKCGLC 133
      || | : | | | :||| | | || |
Db     155 CSECGKTFSRKDNLTQHKRIHTGEMPYKCNCEGKYFSHHSNLIVHQRVHNGARPYKCSDC 214

Qy     134 QEFF-----GLN--NCSSSNRLDG-----HQRVH 155
      : | | :| : | | :|
Db     215 GKVFRHKSTLVQHESIHTGENPYDCSDCGKSGHGYTLIKHQRH 259
```

RESULT 8

T12527

hypothetical protein DKFZp434N043.1 - human (fragment)

C;Species: Homo sapiens (man)

C;Date: 23-Jul-1999 #sequence_revision 23-Jul-1999 #text_change 01-Dec-2000

C;Accession: T12527

R;Wambutt, R.; Heubner, D.; Mewes, H.W.; Gassenhuber, J.; Wiemann, S.

submitted to the Protein Sequence Database, June 1999

A;Reference number: Z17524

A;Accession: T12527

A;Status: preliminary

A;Molecule type: mRNA

A;Residues: 1-386 <WAM>

A;Cross-references: EMBL:AL080143

A;Experimental source: adult testis; clone DKFZp434N043

C;Genetics:

A;Note: DKFZp434N043.1

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 15.4%; Score 136.5; DB 2; Length 386;
Best Local Similarity 20.2%; Pred. No. 1.5e-05;
Matches 47; Conservative 22; Mismatches 73; Indels 91; Gaps 7;

```
Qy      6 AALSCLMLVTALGSMQYQVSSPIYDINYYTSEPC-----QKINVKQ 46
      : | | | : || | | | | : : :
Db     43 STLKCHESVHAREKQGFFVSGKILDQNPQKEKCFKCNKCEKTFSCSKYLTQHERIHTRG 102

Qy     47 IAAYKCGLCAAAQWDFGNT--MCQHQRVHGH-----HHH 78
      : :| | | : : |||:| |
Db    103 VKPFECDCQCGKA---FGQSTRLIHHQRHSRVRLYKWGEQGKAISSASLIKLSFHTKEH 159

Qy     79 SYKCGLCSTRSQKEGLHYT-----CSSHFPYSQY----- 106
      :|| | : : | : | | |
Db    160 PFKCNCEGKTFSHSAHLSKHQLIHAGENPFKCSKCDRVFTQARNYLVQHERTHARKKPLVC 219

Qy    107 ----QFWKNFQTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
      : : | |||:| | | | : ||| | | |||:|
Db    220 NECGKTFRQSSCLSKHQRHSGEKPYVCDYCGKAFL----SAELVRHQRH 267
```

RESULT 9

I39311

Kruppel-type zinc finger protein ZNF74 - human

C;Species: Homo sapiens (man)

C;Date: 06-Sep-1996 #sequence_revision 06-Sep-1996 #text_change 01-Dec-2000

C;Accession: I39311; F42825; S37482

R;Aubry, M.; Demczuk, S.; Desmaze, C.; Aikem, M.; Aurias, A.; Julien, J.P.; Rouleau, G.A.

Hum. Mol. Genet. 2, 1583-1587, 1993

A;Title: Isolation of a zinc finger gene consistently deleted in DiGeorge syndrome.

A;Reference number: I39311; MUID:94093543; PMID:8268910

A;Accession: I39311

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-572 <RES>

A;Cross-references: EMBL:X71623; NID:g1050529; PIDN:CAA50632.1; PID:g1050530

A;Note: submitted to the EMBL Data Library, April 1993, revised 01-NOV-1995

R;Aubry, M.; Marineau, C.; Zhang, F.R.; Zahed, L.; Figlewicz, D.; Delattre, O.; Thomas, G.; de Jong, P.J.; Julien, J.P.; Rouleau, G.A.

Genomics 13, 641-648, 1992

A;Title: Cloning of six new genes with zinc finger motifs mapping to short and long arms of human acrocentric chromosome 22 (p and q11.2).

A;Reference number: A42825; MUID:92347859; PMID:1639391

A;Accession: F42825

A;Status: preliminary; not compared with conceptual translation

A;Molecule type: nucleic acid

A;Residues: 427-444,'V',446-511 <AUB>

A;Note: sequence extracted from NCBI backbone (NCBIP:109775)

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 15.3%; Score 135.5; DB 2; Length 572;

Best Local Similarity 29.2%; Pred. No. 2.7e-05;

Matches 33; Conservative 14; Mismatches 25; Indels 41; Gaps 5;

```

Qy      50 YKCGLCAAAQWDFGNTMCQHQRVHGHHH-----SYKCGLCTRSQKEGLHYTCSSHF 102
          |:||| | | :||| || |:||| | :|||
Db      288 YRCGECGKA-----FNQRTHLTRHHRIHTGEKPYQCGSC-----GKAFTCHS--- 329

Qy      103 YSQYQFWKNFQTLKIHQRVHGSGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
          :|:|::| | :|| |::| | :|:| | |||
Db      330 -----SLTVHEKIHSGDKPFKCSDEKAF-----NSRSRLTLHQRT 366

```

RESULT 10

I48668

zinc finger protein 51 - mouse

N;Alternate names: finger protein zfec12

C;Species: Mus musculus (house mouse)

C;Date: 02-Jul-1996 #sequence_revision 02-Jul-1996 #text_change 08-Dec-2000

C;Accession: I48668; A40984

R;Burke, P.S.; Don, J.; Wolgemuth, D.J.

Mamm. Genome 5, 387-389, 1994

A;Title: Zfp-51, a murine zinc finger encoding gene mapping to the t-complex region of chromosome 17, encodes 19 contiguous zinc fingers and is ubiquitously expressed.

A;Reference number: I48668; MUID:94319090; PMID:8043957

A;Accession: I48668

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-710 <RES>

A;Cross-references: EMBL:X74855; NID:g488832; PIDN:CAA52847.1; PID:g488833

R;Crossley, P.H.; Little, P.F.R.

Proc. Natl. Acad. Sci. U.S.A. 88, 7923-7927, 1991

A;Title: A cluster of related zinc finger protein genes is deleted in the mouse embryonic lethal mutation t(w18).

A;Reference number: A40984; MUID:91376058; PMID:1680234

A;Accession: A40984

A;Status: preliminary

A;Molecule type: mRNA

A;Residues: 672-710 <CRO>

A;Cross-references: GB:M74235

C;Genetics:

A;Gene: Zfp-51

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

C;Keywords: DNA binding; zinc finger

Query Match 15.2%; Score 134; DB 2; Length 710;
Best Local Similarity 28.6%; Pred. No. 4.7e-05;
Matches 38; Conservative 20; Mismatches 39; Indels 36; Gaps 7;

```
Qy      28 IYDINYYS-----EPCQKINVKQIAAYKCGLCAAAQWDFGNTMCQHQRVHGHHSYKC 82
      :|| :|::|      :|  |  |  ||| |      :: ::| |::|  |||
Db      120 VYD-DYFSSAYSILMQPTAYIKEKPHQCCKGCGKC----FNTSSSLTVHHRH-TKEKPYKC 173

Qy      83 GLCTRSQKEGLHYTCSSHFYPYSQYQFWKNFQTLKIHQRVHGGGGSYKCGLCQEFFGLNNC 142
      :| :|      :| :|      ||||| | |  ||| |::|
Db      174 SVCDKS-----FTQCTH-----LKIHQRRHTGEKPYKCKECEKSF----- 208

Qy      143 SSSNRLDGHQRVH 155
      : |  ||::|
Db      209 VQLSALKSHQKLH 221
```

RESULT 11

S03679

finger protein (clone mkr5) - mouse (fragment)

C;Species: Mus musculus (house mouse)

C;Date: 04-Dec-1992 #sequence_revision 04-Dec-1992 #text_change 01-Dec-2000

C;Accession: S03679

R;Chowdhury, K.; Rohdewohld, H.; Gruss, P.

Nucleic Acids Res. 16, 9995-10011, 1988

A;Title: Specific and ubiquitous expression of different Zn finger protein genes in the mouse.

A;Reference number: S03677; MUID:89057528; PMID:3143103

A;Accession: S03679

A;Status: not compared with conceptual translation

A;Molecule type: mRNA

A;Residues: 1-475 <CHO>

A;Note: the sequence in fig.2 is inconsistent with that shown in fig.1 in having 64-Asn, 159-Trp, 187-trp, 188-trp, and in lacking a Met after 64-Gly and an Asp after 78-Ser

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 15.1%; Score 133.5; DB 2; Length 475;
Best Local Similarity 27.8%; Pred. No. 3.5e-05;
Matches 40; Conservative 24; Mismatches 49; Indels 31; Gaps 8;

```
Qy      34 YTSEPCQK-----INVKQ-----IAAYKCGLCAAAQWDFGNTMCQHQRVHGHHSYK 81
      || || | : | | | : | : | : | : | : | : |
Db      223 YTCEVCHKSFYRGSSLTVHQRIHTGEKPYECEICRKA-FSHHASLTQHQRVHS-GEKPFK 280

Qy      82 CGLCTRSQKEGLHYTCSSHF-----PYSQYQFWKNF---QTLKIHQRVHGGGGSYKCG 131
      | | : : : : | : | : | : | : | : | : | : |
Db      281 CKECGKAFRQNIH--LASHWRIHTGEKPFECGCGKSFSISSQLATHQRIHTGEKPFECK 338

Qy     132 LCQEFFGLNNCSSSNRLDGHQRVH 155
      :| : | | | | | | : |
Db     339 VCRKAFRQN-----IHLASHWRIH 357
```

RESULT 12

I48689

gene NK10 protein - mouse

C;Species: Mus musculus (house mouse)

C;Date: 02-Jul-1996 #sequence_revision 02-Jul-1996 #text_change 01-Dec-2000

C;Accession: I48689; S49078

R;Lange, R.; Christoph, A.; Thiesen, H.J.; Vopper, G.; Johnson, K.R.; Lemaire, L.; Plomann, M.; Cremer, H.; Barthels, D.; Heinleim, U.A.O.

DNA Cell Biol. 14, 971-981, 1995

A;Title: Developmentally regulated mouse gene NK10 eucodes a xZinc Finger Repressor Protein with differential NDA-Binding Domains.

A;Reference number: I48689; MUID:96069544; PMID:7576184

A;Accession: I48689

A;Status: preliminary; translated from GB/EMBL/DDBJ

A;Molecule type: mRNA

A;Residues: 1-636 <RES>

A;Cross-references: EMBL:X79828; NID:g506501; PIDN:CAA56225.1; PID:g506502

C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 15.1%; Score 133.5; DB 2; Length 636;
Best Local Similarity 31.6%; Pred. No. 4.7e-05;
Matches 37; Conservative 15; Mismatches 44; Indels 21; Gaps 6;

```
Qy      50 YKCGLCAAAQWDFGNTMCQHQRVHGHHSYKCGLCCTRS-----QKEGLHYTCSSHF 101
      |:| || | : : : | || : | :| || | | | | | |
Db      306 YQCSLCGKA-FQRSSSLVQHQRH-TGEKPYRCNLGSRFRHSTSLTQHEVTH---SGEK 360

Qy     102 PYSQYQFWKNF---QTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
      |: : | | : | :| | | :| :| | | | | | |
Db     361 PFQCKEKGKAFSRCSSSLVQHERHTTGEKPFECGICGRAFG-----QSPSLYKHMRIH 412
```

RESULT 13

T14757

hypothetical protein DKFZp572C163.1 - human (fragment)

C;Species: Homo sapiens (man)
 C;Date: 20-Sep-1999 #sequence_revision 20-Sep-1999 #text_change 01-Dec-2000
 C;Accession: T14757
 R;Wambutt, R.; Heubner, D.; Mewes, H.W.; Gassenhuber, J.; Wiemann, S.
 submitted to the Protein Sequence Database, August 1999
 A;Reference number: Z18181
 A;Accession: T14757
 A;Status: preliminary
 A;Molecule type: mRNA
 A;Residues: 1-701 <WAM>
 A;Cross-references: EMBL:AL110217
 A;Experimental source: adult subthalamic nucleus; clone DKFZp572C163
 C;Genetics:
 A;Note: DKFZp572C163.1
 C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology

Query Match 15.1%; Score 133.5; DB 2; Length 701;
 Best Local Similarity 23.4%; Pred. No. 5.2e-05;
 Matches 41; Conservative 16; Mismatches 37; Indels 81; Gaps 8;

```

Qy      50 YKCGLCAAAQWDFGNT-----MCQHQRVH-----GHHH-HS- 79
          |:| | | | | | | | | | | | | | | | | | | | | | | | | | | |
Db      468 YECNQC-----GKTFSQRTHLCAHQRIHTGEKPYECNECGKTFADNSALRAHHRIHTG 520

Qy      80 ---YKCGLCTRSQKEGLHY-----TCSSHFPYSQYQFWKNF----- 112
          |:| | : : | | | | | | | | | | | | | | | | | | | | | |
Db      521 EKPYECDNCGKTFSKTSHLRAHLRTRSGEKPYECSECGKTFSEKSYVSAHQRVHTGEKPY 580

Qy      113 -----QTLKIHQRVHGGGGSYKCGLCQEFFGLNNCSSSNRLDGHQRVH 155
          ||:| |||:| | ||:| | : | | | : | | ||:|
Db      581 ECNVCGKPFahnSTLRVHQRIHTGEKSYECNDCGKTF-----SQKSHLSAHQRIH 630

```

RESULT 14

A34612

zinc finger protein ZNF7 - human

N;Alternate names: zinc finger protein kox4

C;Species: Homo sapiens (man)

C;Date: 22-Jun-1990 #sequence_revision 22-Jun-1990 #text_change 01-Dec-2000

C;Accession: A34612; A56409; S10421; I37972

R;Lania, L.; Donti, E.; Pannuti, A.; Pascucci, A.; Pengue, G.; Feliciello, I.;

La Mantia, G.; Lanfranccone, L.; Pelicci, P.G.

Genomics 6, 333-340, 1990

A;Title: cDNA isolation, expression analysis, and chromosomal localization of two human zinc finger genes.

A;Reference number: A34612; MUID:90169993; PMID:2106481

A;Accession: A34612

A;Molecule type: mRNA

A;Residues: 1-686 <LAN>

A;Cross-references: GB:M29580; NID:g340445; PIDN:AAA61313.1; PID:g340446; GB:J04751

R;Bray, P.; Lichter, P.; Thiesen, H.J.; Ward, D.C.; Dawid, I.B.

Proc. Natl. Acad. Sci. U.S.A. 88, 9563-9567, 1991

A;Title: Characterization and mapping of human genes encoding zinc finger proteins.

A;Reference number: A56409; MUID:92052132; PMID:1946370

A;Accession: A56409

A;Molecule type: DNA
 A;Residues: 425-589 <BRA>
 A;Cross-references: GB:M77170
 R;Thiesen, H.J.
 submitted to the EMBL Data Library, March 1990
 A;Reference number: S10397
 A;Accession: S10421
 A;Molecule type: mRNA
 A;Residues: 413-468 <THI>
 A;Cross-references: EMBL:X52335; NID:g34165; PIDN:CAA36561.1; PID:g930095
 R;Thiesen, H.J.
 New Biol. 2, 363-374, 1990
 A;Title: Multiple genes encoding zinc finger domains are expressed in human T cells.
 A;Reference number: I37949; MUID:91145339; PMID:2288909
 A;Accession: I37972
 A;Status: translated from GB/EMBL/DDBJ
 A;Molecule type: mRNA
 A;Residues: 413-468 <RES>
 A;Cross-references: EMBL:X52335; NID:g34165; PIDN:CAA36561.1; PID:g930095
 C;Genetics:
 A;Gene: GDB:ZNF7
 A;Cross-references: GDB:120509; OMIM:194531
 A;Map position: 8q24.3-8q24.3
 C;Superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat homology
 C;Keywords: DNA binding; zinc finger

Query Match 14.9%; Score 131.5; DB 2; Length 686;
 Best Local Similarity 26.9%; Pred. No. 7.9e-05;
 Matches 45; Conservative 11; Mismatches 46; Indels 65; Gaps 8;

```

Qy      50 YKCGLCAAAQWDFG--NTMCQHQRVHGHHSYKCGLCTRSQKEGLH----- 94
      ||| | | || : : :||| | :|| | : :| |
Db      441 YKCNKCTKA---FGCSSRLIRHQRTHTGEKPFKCECGKGFVQGSGLIQQHQRIHTGEKP 496

Qy      95 YTCS-----SHFPYSQYQFWKNFQ---TLKIHQRVHGGGGSYKCG 131
      | | :                || | | | | | | | | | | |
Db      497 YVCNDCGKAQFSQSSSLIYHQRIHKGEKPYECLQCGKAQFSMSTQLTIHQRVHTGERPYKCN 556

Qy      132 LCQEFFGLNN-----CSS-----SNRLDGHQRVH 155
      | : | | :                || | : | | | : |
Db      557 ECGKAQFSQNSTLFQHQIIHAGVKPYECSECGKAQFSRSSYLIHQRIH 603
  
```

RESULT 15

S70006

finger protein zFOC1 - guinea pig

C;Species: Cavia porcellus (guinea pig)

C;Date: 06-Dec-1996 #sequence_revision 13-Mar-1997 #text_change 05-Nov-1999

C;Accession: S70006

R;Rivolta, M.N.; Negrini, C.; Wilcox, E.R.

Biochim. Biophys. Acta 1306, 127-132, 1996

A;Title: A novel zinc finger gene preferentially expressed in the retina and the organ of Corti localizes to human chromosome 12q24.3.

A;Reference number: S70006; MUID:96221281; PMID:8634327

A;Accession: S70006

A;Status: preliminary

A;Molecule type: mRNA
 A;Residues: 1-261 <RIV>
 A;Cross-references: EMBL:L26335; NID:g1237277; PIDN:AAC42091.1; PID:g1237278
 C;Keywords: zinc

Query Match 14.8%; Score 131; DB 2; Length 261;
 Best Local Similarity 28.8%; Pred. No. 3.4e-05;
 Matches 42; Conservative 16; Mismatches 44; Indels 44; Gaps 7;

```

Qy      45 KQIAAYKCG-----LCAAAQWDFGNT--MCQHQRVHGHHHSY 80
      |   |::|               :|:   | |:  :| |||||   :
Db      85 KPYKCYECGKAFNWSPHLQIHMRVHTGEKPYVCSECGRGFSNSSNLCMHQRVH-TGEKPF 143

Qy      81 KCGLCTRS-----QKEGLHYTCSSHFYPYSQYQFWKNF---QTLKIHQRVHGGGGSYK 129
      ||  | ::           : :|   :  ||  |:  ||   :| ||||| |  |:
Db     144 KCEECGKAfrHTSSSLCMHQRVH---TGEKPYKCYECGKAfSQSSSLCIHQrvHTGEKPYR 200

Qy     130 CGLCQEFFGLNNCSSSNRLDGHQRVH 155
      |  | : |   | |: |  |||||
Db     201 CCGCGKAF-----SQSSSLCIHQrvH 221
  
```

Search completed: March 5, 2004, 16:28:57
 Job time : 35.858 secs

GenCore version 5.1.6
Copyright (c) 1993 - 2004 Compugen Ltd.

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:22:54 ; Search time 84.7994 Seconds
(without alignments)
390.935 Million cell updates/sec

Title: US-10-057-890A-31
Perfect score: 884
Sequence: 1 MKVSVAALSCLMLVTALGSM.....GLNNCSSSNRLDGHQRVHAA 157

Scoring table: BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 809742 seqs, 211153259 residues

Total number of hits satisfying chosen parameters: 809742

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : Published Applications_AA:*

- 1: /cgn2_6/ptodata/1/pubpaa/US07_PUBCOMB.pep:*
- 2: /cgn2_6/ptodata/1/pubpaa/PCT_NEW_PUB.pep:*
- 3: /cgn2_6/ptodata/1/pubpaa/US06_NEW_PUB.pep:*
- 4: /cgn2_6/ptodata/1/pubpaa/US06_PUBCOMB.pep:*
- 5: /cgn2_6/ptodata/1/pubpaa/US07_NEW_PUB.pep:*
- 6: /cgn2_6/ptodata/1/pubpaa/PCTUS_PUBCOMB.pep:*
- 7: /cgn2_6/ptodata/1/pubpaa/US08_NEW_PUB.pep:*
- 8: /cgn2_6/ptodata/1/pubpaa/US08_PUBCOMB.pep:*
- 9: /cgn2_6/ptodata/1/pubpaa/US09A_PUBCOMB.pep:*
- 10: /cgn2_6/ptodata/1/pubpaa/US09B_PUBCOMB.pep:*
- 11: /cgn2_6/ptodata/1/pubpaa/US09C_PUBCOMB.pep:*
- 12: /cgn2_6/ptodata/1/pubpaa/US09_NEW_PUB.pep:*
- 13: /cgn2_6/ptodata/1/pubpaa/US10A_PUBCOMB.pep:*
- 14: /cgn2_6/ptodata/1/pubpaa/US10B_PUBCOMB.pep:*
- 15: /cgn2_6/ptodata/1/pubpaa/US10C_PUBCOMB.pep:*
- 16: /cgn2_6/ptodata/1/pubpaa/US10_NEW_PUB.pep:*
- 17: /cgn2_6/ptodata/1/pubpaa/US60_NEW_PUB.pep:*
- 18: /cgn2_6/ptodata/1/pubpaa/US60_PUBCOMB.pep:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result	%	Query				
No.	Score	Match Length DB ID				Description

1	884	100.0	157	14	US-10-057-890A-31	Sequence 31, Appl
2	797	90.2	138	14	US-10-057-890A-10	Sequence 10, Appl
3	379	42.9	352	9	US-09-725-285-2	Sequence 2, Appli
4	379	42.9	352	9	US-09-759-841-2	Sequence 2, Appli
5	379	42.9	352	9	US-09-779-879A-22	Sequence 22, Appl
6	379	42.9	352	9	US-09-779-880A-22	Sequence 22, Appl
7	379	42.9	352	9	US-09-813-653-15	Sequence 15, Appl
8	379	42.9	352	9	US-09-813-653-17	Sequence 17, Appl
9	379	42.9	352	9	US-09-796-202-1	Sequence 1, Appli
10	379	42.9	352	9	US-09-195-662A-2	Sequence 2, Appli
11	379	42.9	352	9	US-09-339-912A-2	Sequence 2, Appli
12	379	42.9	352	9	US-09-938-719-5	Sequence 5, Appli
13	379	42.9	352	9	US-09-939-226-5	Sequence 5, Appli
14	379	42.9	352	9	US-09-938-703-5	Sequence 5, Appli
15	379	42.9	352	9	US-09-502-783A-2	Sequence 2, Appli
16	379	42.9	352	10	US-09-734-221A-14	Sequence 14, Appl
17	379	42.9	352	11	US-09-826-509-477	Sequence 477, App
18	379	42.9	352	13	US-10-106-623-2	Sequence 2, Appli
19	379	42.9	352	14	US-10-232-686-2	Sequence 2, Appli
20	379	42.9	352	14	US-10-086-814-1	Sequence 1, Appli
21	379	42.9	352	14	US-10-067-800-22	Sequence 22, Appl
22	379	42.9	352	14	US-10-290-058A-6	Sequence 6, Appli
23	379	42.9	352	14	US-10-225-567A-352	Sequence 352, App
24	379	42.9	352	14	US-10-323-314-1	Sequence 1, Appli
25	379	42.9	352	14	US-10-072-301-1	Sequence 1, Appli
26	379	42.9	352	14	US-10-071-866-1	Sequence 1, Appli
27	379	42.9	352	14	US-10-135-839-22	Sequence 22, Appl
28	379	42.9	352	14	US-10-239-423-67	Sequence 67, Appl
29	379	42.9	352	14	US-10-439-845-4	Sequence 4, Appli
30	379	42.9	352	15	US-10-360-828-1	Sequence 1, Appli
31	374	42.3	352	14	US-10-164-649-52	Sequence 52, Appl
32	374	42.3	352	14	US-10-439-845-2	Sequence 2, Appli
33	373	42.2	352	9	US-09-779-879A-2	Sequence 2, Appli
34	373	42.2	352	9	US-09-779-880A-2	Sequence 2, Appli
35	373	42.2	352	14	US-10-067-800-2	Sequence 2, Appli
36	373	42.2	352	14	US-10-135-839-2	Sequence 2, Appli
37	363	41.1	352	13	US-10-106-623-20	Sequence 20, Appl
38	258	29.2	184	9	US-09-938-719-4	Sequence 4, Appli
39	258	29.2	184	9	US-09-939-226-4	Sequence 4, Appli
40	258	29.2	184	9	US-09-938-703-4	Sequence 4, Appli
41	258	29.2	215	9	US-09-938-719-6	Sequence 6, Appli
42	258	29.2	215	9	US-09-939-226-6	Sequence 6, Appli
43	258	29.2	215	9	US-09-938-703-6	Sequence 6, Appli
44	183.5	20.8	332	14	US-10-095-876A-2	Sequence 2, Appli
45	182	20.6	32	14	US-10-057-890A-13	Sequence 13, Appl

ALIGNMENTS

RESULT 1
 US-10-057-890A-31
 ; Sequence 31, Application US/10057890A
 ; Publication No. US20030044901A1
 ; GENERAL INFORMATION:
 ; APPLICANT: Coleman, Timothy

```
; APPLICANT: Mansfield, Brian
; TITLE OF INVENTION: Scaffold Fusion Polypeptides, Composition for Making the
Same, and Methods
; TITLE OF INVENTION: of Using the Same.
; FILE REFERENCE: PF537
; CURRENT APPLICATION NUMBER: US/10/057,890A
; CURRENT FILING DATE: 2002-01-29
; PRIOR APPLICATION NUMBER: 60/265,782
; PRIOR FILING DATE: 2001-01-31
; PRIOR APPLICATION NUMBER: 60/265,858
; PRIOR FILING DATE: 2001-01-31
; NUMBER OF SEQ ID NOS: 32
; SEQ ID NO 31
; LENGTH: 157
; TYPE: PRT
; ORGANISM: Homo sapiens
US-10-057-890A-31
```

```
Query Match          100.0%; Score 884; DB 14; Length 157;
Best Local Similarity 100.0%; Pred. No. 7.8e-80;
Matches 157; Conservative 0; Mismatches 0; Indels 0; Gaps 0;
```

```
Qy      1 MKVSVAALSCLMLVTALGSMDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAQW 60
          |||
Db      1 MKVSVAALSCLMLVTALGSMDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAQW 60

Qy      61 DFGNTMCQHQRVHGHHSYKCGCLTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQR 120
          |||
Db      61 DFGNTMCQHQRVHGHHSYKCGCLTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQR 120

Qy      121 VHGGGGSYKCGLCQEFFGLNNCSSSRLDGHQRVHAA 157
          |||
Db      121 VHGGGGSYKCGLCQEFFGLNNCSSSRLDGHQRVHAA 157
```

RESULT 2

US-10-057-890A-10

```
; Sequence 10, Application US/10057890A
; Publication No. US20030044901A1
; GENERAL INFORMATION:
; APPLICANT: Coleman, Timothy
; APPLICANT: Mansfield, Brian
; TITLE OF INVENTION: Scaffold Fusion Polypeptides, Composition for Making the
Same, and Methods
; TITLE OF INVENTION: of Using the Same.
; FILE REFERENCE: PF537
; CURRENT APPLICATION NUMBER: US/10/057,890A
; CURRENT FILING DATE: 2002-01-29
; PRIOR APPLICATION NUMBER: 60/265,782
; PRIOR FILING DATE: 2001-01-31
; PRIOR APPLICATION NUMBER: 60/265,858
; PRIOR FILING DATE: 2001-01-31
; NUMBER OF SEQ ID NOS: 32
; SEQ ID NO 10
; LENGTH: 138
; TYPE: PRT
; ORGANISM: Homo sapiens
```

US-10-057-890A-10

Query Match 90.2%; Score 797; DB 14; Length 138;
Best Local Similarity 100.0%; Pred. No. 2.8e-71;
Matches 138; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAAQWDFGNTMCQHQRVHGHHHHS 79
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAAYKCGLCAAAQWDFGNTMCQHQRVHGHHHHS 60

Qy      80 YKCGLCTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQRVHGGSYKCGLCQEFFGL 139
          |||
Db      61 YKCGLCTRSQKEGLHYTCSSHPYSQYQFWKNFQTLKIHQRVHGGSYKCGLCQEFFGL 120

Qy      140 NNCSSSNRLDGHQRVHAA 157
          |||
Db      121 NNCSSSNRLDGHQRVHAA 138
```

RESULT 3

US-09-725-285-2

```
; Sequence 2, Application US/09725285
; Patent No. US20010000241A1
; GENERAL INFORMATION:
; APPLICANT: Li, Yi
; APPLICANT: Ruben, Steven, M.
; TITLE OF INVENTION: Antibodies to Human G-Protein Chemokine Receptor
HDGNR10
; TITLE OF INVENTION: (CCR5 Receptor)
; FILE REFERENCE: 1488.1150003
; CURRENT APPLICATION NUMBER: US/09/725,285
; CURRENT FILING DATE: 2000-11-29
; PRIOR APPLICATION NUMBER: 09/339,912
; PRIOR FILING DATE: 1999-06-25
; PRIOR APPLICATION NUMBER: 09/195,662
; PRIOR FILING DATE: 1998-11-18
; PRIOR APPLICATION NUMBER: 08/466,343
; PRIOR FILING DATE: 1995-06-06
; NUMBER OF SEQ ID NOS: 9
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 2
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Artificial Sequence: Genomic
; FEATURE:
; OTHER INFORMATION: Deduced Amino Acid Sequence
US-09-725-285-2
```

Query Match 42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHHSYKCG--- 83
```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPLLYSLVFIFGFVGNMLVILILINCKR 60

Qy     50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHHSYKCG--- 83
      |  |                               |||||  | : | : |
Db     61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIFGFFSGIFF 118

Qy     84 -----LCTRSQKEGLHYTC 97
      : ||||||||||||
Db    119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSOKEGLHYTC 178

```



```

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      |||
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
      | | |||
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 5

```

US-09-779-879A-22
; Sequence 22, Application US/09779879A
; Patent No. US20020048786A1
; GENERAL INFORMATION:
; APPLICANT: Rosen, Craig A.
; APPLICANT: Roschke, Viktor
; APPLICANT: Li, Yi
; APPLICANT: Ruben, Steven, M.
; TITLE OF INVENTION: Human G-protein Chemokine Receptor (CCR5) HDGNR10
; FILE REFERENCE: 1488.115000A
; CURRENT APPLICATION NUMBER: US/09/779,879A
; CURRENT FILING DATE: 2001-02-09
; PRIOR APPLICATION NUMBER: US 60/181,258
; PRIOR FILING DATE: 2000-02-09
; PRIOR APPLICATION NUMBER: US 60/187,999
; PRIOR FILING DATE: 2000-03-09
; PRIOR APPLICATION NUMBER: US 60/234,336
; PRIOR FILING DATE: 2000-09-22
; NUMBER OF SEQ ID NOS: 58
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 22
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Homo sapiens
US-09-779-879A-22

```

```

Query Match          42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | |||
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : |||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      |||
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238

```

Qy 125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
| | | | | | | | | | | | | | | | | | : |
Db 239 TIMIVYFLFWAPYNIVLLNTFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 6

US-09-779-880A-22
; Sequence 22, Application US/09779880A
; Patent No. US20020061834A1
; GENERAL INFORMATION:
; APPLICANT: Rosen, Craig A.
; APPLICANT: Roschke, Viktor
; APPLICANT: Li, Yi
; APPLICANT: Ruben, Steven, M.
; TITLE OF INVENTION: Human G-protein Chemokine Receptor (CCR5) HDGNR10
; FILE REFERENCE: 1488.115000C
; CURRENT APPLICATION NUMBER: US/09/779,880A
; CURRENT FILING DATE: 2001-02-09
; PRIOR APPLICATION NUMBER: US 60/181,258
; PRIOR FILING DATE: 2000-02-09
; PRIOR APPLICATION NUMBER: US 60/187,999
; PRIOR FILING DATE: 2000-03-09
; PRIOR APPLICATION NUMBER: US 60/234,336
; PRIOR FILING DATE: 2000-09-22
; NUMBER OF SEQ ID NOS: 58
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 22
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Homo sapiens
US-09-779-880A-22

Query Match 42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
| | | | | | | | | | | | | | | | | |
Db 1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
| | | | | | | | | | | | | | | | : | : : |
Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy 84 -----LCTRSQKEGLHYTC 97
: | | | | | | | | | |
Db 119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
| | | | | | | | | | | | | | | | : : |
Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy 125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
| | | | | | | | | | | | | | | | : |
Db 239 TIMIVYFLFWAPYNIVLLNTFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 7

US-09-813-653-15

; Sequence 15, Application US/09813653

; Patent No. US20020064770A1

; GENERAL INFORMATION:

; APPLICANT: Nestor, John

; APPLICANT: Wilson, Carol

; APPLICANT: See, Raymond

; APPLICANT: Tan Hehir, Christina

; TITLE OF INVENTION: Binding Compounds and Methods For Identifying Binding Compounds

; FILE REFERENCE: CNS-005

; CURRENT APPLICATION NUMBER: US/09/813,653

; CURRENT FILING DATE: 2001-03-20

; PRIOR APPLICATION NUMBER: US 60/190,946

; PRIOR FILING DATE: 2000-03-21

; PRIOR APPLICATION NUMBER: US 60/190,996

; PRIOR FILING DATE: 2000-03-21

; PRIOR APPLICATION NUMBER: US 60/191,299

; PRIOR FILING DATE: 2000-03-21

; NUMBER OF SEQ ID NOS: 44

; SOFTWARE: PatentIn version 3.0

; SEQ ID NO 15

; LENGTH: 352

; TYPE: PRT

; ORGANISM: Homo sapiens

US-09-813-653-15

Query Match 42.9%; Score 379; DB 9; Length 352;

Best Local Similarity 35.0%; Pred. No. 1.8e-29;

Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
          |  |
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFGFLNNCSSSNRLDQAMQV 281

```

RESULT 8

US-09-813-653-17

; Sequence 17, Application US/09813653

; Patent No. US20020064770A1

```

; GENERAL INFORMATION:
; APPLICANT: Nestor, John
; APPLICANT: Wilson, Carol
; APPLICANT: See, Raymond
; APPLICANT: Tan Hehir, Christina
; TITLE OF INVENTION: Binding Compounds and Methods For Identifying Binding
Compounds
; FILE REFERENCE: CNS-005
; CURRENT APPLICATION NUMBER: US/09/813,653
; CURRENT FILING DATE: 2001-03-20
; PRIOR APPLICATION NUMBER: US 60/190,946
; PRIOR FILING DATE: 2000-03-21
; PRIOR APPLICATION NUMBER: US 60/190,996
; PRIOR FILING DATE: 2000-03-21
; PRIOR APPLICATION NUMBER: US 60/191,299
; PRIOR FILING DATE: 2000-03-21
; NUMBER OF SEQ ID NOS: 44
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 17
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Homo sapiens
US-09-813-653-17

```

```

Query Match          42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILIQINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               |||||   | : | :   : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||||           : : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLPPLVMVICYSGLKTLRLCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC-----QEFGGLNNCSSSNRLDGHQRV 154
          |   |   |||||||||||||   :|
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFGGLNNCSSSNRLDQAMQV 281

```

RESULT 9

US-09-796-202-1

; Sequence 1, Application US/09796202

; Patent No. US20020068813A1

; GENERAL INFORMATION:

; APPLICANT: Dragic, Tatjana

; APPLICANT: Olson, William

; TITLE OF INVENTION: SULFATED CCR5 PEPTIDES FOR HIV-1 INFECTION

; FILE REFERENCE: 2048/61010/JPW/SHS
; CURRENT APPLICATION NUMBER: US/09/796,202
; CURRENT FILING DATE: 2001-02-28
; NUMBER OF SEQ ID NOS: 17
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 1
; LENGTH: 352
; TYPE: PRT
; ORGANISM: human
US-09-796-202-1

Query Match 42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGLIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC---QEFFGLNNC SSSNRLDGHQRV 154
          |  |
Db     239 TIMIVYFLEWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281
```

RESULT 10

US-09-195-662A-2

; Sequence 2, Application US/09195662A
; Patent No. US20020076745A1
; GENERAL INFORMATION:
; APPLICANT: Li, Yi
; APPLICANT: Ruben, Steven, M.
; TITLE OF INVENTION: Human G-Protein Chemokine Receptor HDG NR10 (CCR5 Receptor)
; FILE REFERENCE: 1488.1150002
; CURRENT APPLICATION NUMBER: US/09/195,662A
; CURRENT FILING DATE: 1998-11-18
; PRIOR APPLICATION NUMBER: 08/466,343
; PRIOR FILING DATE: 1995-06-06
; NUMBER OF SEQ ID NOS: 9
; SOFTWARE: PatentIn version 3.0
; SEQ ID NO 2
; LENGTH: 352
; TYPE: PRT
; ORGANISM: Artificial Sequence: Genomic
; FEATURE:

; OTHER INFORMATION: Deduced Amino Acid Sequence
US-09-195-662A-2

Query Match 42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;
Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||||||  :: |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
Db     239 TIMIVYFLEWAPYNIVLLNLNTFQEFFGLNNCSSSNRLDQAMQV 281
```

RESULT 11

US-09-339-912A-2

; Sequence 2, Application US/09339912A

; Patent No. US20020099176A1

; GENERAL INFORMATION:

; APPLICANT: Li, Yi

; APPLICANT: Ruben, Steven, M.

; TITLE OF INVENTION: Antibodies to Human G-Protein Chemokine Receptor
HDG NR10

; TITLE OF INVENTION: (CCR5 Receptor)

; FILE REFERENCE: 1488.1150003

; CURRENT APPLICATION NUMBER: US/09/339,912A

; CURRENT FILING DATE: 1999-06-25

; PRIOR APPLICATION NUMBER: 09/195,662

; PRIOR FILING DATE: 1998-11-18

; PRIOR APPLICATION NUMBER: 08/466,343

; PRIOR FILING DATE: 1995-06-06

; NUMBER OF SEQ ID NOS: 9

; SOFTWARE: PatentIn version 3.0

; SEQ ID NO 2

; LENGTH: 352

; TYPE: PRT

; ORGANISM: Artificial Sequence: Genomic

; FEATURE:

; OTHER INFORMATION: Deduced Amino Acid Sequence

US-09-339-912A-2

Query Match 42.9%; Score 379; DB 9; Length 352;
Best Local Similarity 35.0%; Pred. No. 1.8e-29;

Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      ||||||||||||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
      | | |||||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : ||||||||
Db     119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      |||||||||||||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
      | | |||||||||||| : |
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 12

US-09-938-719-5

; Sequence 5, Application US/09938719

; Patent No. US20020106742A1

; GENERAL INFORMATION:

; APPLICANT: SAMSON, MICHEL

; PARMENTIER, MARC

; VASSART, GILBERT

; LIBERT, FREDERICK

; TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR

; AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR

; NUMBER OF SEQUENCES: 17

; CORRESPONDENCE ADDRESS:

; ADDRESSEE: Knobbe, Martens, Olson & Bear

; STREET: 620 Newport Center Drive 16th Floor

; CITY: Newport Beach

; STATE: CA

; COUNTRY: U.S.A.

; ZIP: 92660

; COMPUTER READABLE FORM:

; MEDIUM TYPE: Floppy disk

; COMPUTER: IBM PC compatible

; OPERATING SYSTEM: PC-DOS/MS-DOS

; SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

; CURRENT APPLICATION DATA:

; APPLICATION NUMBER: US/09/938,719

; FILING DATE: 24-Aug-2001

; CLASSIFICATION: <Unknown>

; PRIOR APPLICATION DATA:

; APPLICATION NUMBER: 09/626,939

; FILING DATE: 27-JULY-2000

; ATTORNEY/AGENT INFORMATION:

; NAME: Altman, Daniel E

; REGISTRATION NUMBER: 34,115

```

;          REFERENCE/DOCKET NUMBER: <Unknown>
;  INFORMATION FOR SEQ ID NO: 5:
;    SEQUENCE CHARACTERISTICS:
;      LENGTH: 352 amino acids
;      TYPE: amino acid
;      TOPOLOGY: linear
;    MOLECULE TYPE: protein
;    SEQUENCE DESCRIPTION: SEQ ID NO: 5:
US-09-938-719-5

```

```

Query Match          42.9%;  Score 379;  DB 9;  Length 352;
Best Local Similarity 35.0%;  Pred. No. 1.8e-29;
Matches 99;  Conservative 7;  Mismatches 27;  Indels 150;  Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  | : | :  |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGLIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||||||||||           :: |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC---QEFFGLNNC SSSNRLDGHQRV 154
          |  |                      |||||  :|
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281

```

RESULT 13

US-09-939-226-5

```

; Sequence 5, Application US/09939226
; Patent No. US20020110805A1
;  GENERAL INFORMATION:
;    APPLICANT: SAMSON, MICHEL
;              PARMENTIER, MARC
;              VASSART, GILBERT
;              LIBERT, FREDERICK
;    TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR
;                      AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR
;    NUMBER OF SEQUENCES: 17
;    CORRESPONDENCE ADDRESS:
;      ADDRESSEE: Knobbe, Martens, Olson & Bear
;      STREET: 620 Newport Center Drive 16th Floor
;      CITY: Newport Beach
;      STATE: CA
;      COUNTRY: U.S.A.
;      ZIP: 92660
;    COMPUTER READABLE FORM:
;      MEDIUM TYPE: Floppy disk
;      COMPUTER: IBM PC compatible

```



```

;           OPERATING SYSTEM: PC-DOS/MS-DOS
;           SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)
;
;   CURRENT APPLICATION DATA:
;           APPLICATION NUMBER: US/09/939,226
;           FILING DATE: 24-Aug-2001
;           CLASSIFICATION: <Unknown>
;
;   PRIOR APPLICATION DATA:
;           APPLICATION NUMBER: 09/626,939
;           FILING DATE: 2000-07-27
;
;   ATTORNEY/AGENT INFORMATION:
;           NAME: Altman, Daniel E
;           REGISTRATION NUMBER: 34,115
;           REFERENCE/DOCKET NUMBER: <Unknown>
;
;   INFORMATION FOR SEQ ID NO: 5:
;           SEQUENCE CHARACTERISTICS:
;               LENGTH: 352 amino acids
;               TYPE: amino acid
;               TOPOLOGY: linear
;           MOLECULE TYPE: protein
;           SEQUENCE DESCRIPTION: SEQ ID NO: 5:
US-09-939-226-5

```

```

Query Match          42.9%;  Score 379;  DB 9;  Length 352;
Best Local Similarity 35.0%;  Pred. No. 1.8e-29;
Matches 99;  Conservative 7;  Mismatches 27;  Indels 150;  Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||||||||||||||||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |  |                      |||||  | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||||  :: |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKCGLC----QEFGGLNNCSSSNRLDGHQRV 154
          |  |  |||||  :|
Db      239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 14

US-09-938-703-5

; Sequence 5, Application US/09938703

; Patent No. US20020110870A1

; GENERAL INFORMATION:

; APPLICANT: SAMSON, MICHEL

; PARMENTIER, MARC

; VASSART, GILBERT

; LIBERT, FREDERICK

; TITLE OF INVENTION: ACTIVE AND INACTIVE CC-CHEMOKINES RECEPTOR

```

;                               AND NUCLEIC ACID MOLECULES ENCODING SAID RECEPTOR
;
;   NUMBER OF SEQUENCES: 17
;   CORRESPONDENCE ADDRESS:
;       ADDRESSEE: Knobbe, Martens, Olson & Bear
;       STREET: 620 Newport Center Drive 16th Floor
;       CITY: Newport Beach
;       STATE: CA
;       COUNTRY: U.S.A.
;       ZIP: 92660
;
;   COMPUTER READABLE FORM:
;       MEDIUM TYPE: Floppy disk
;       COMPUTER: IBM PC compatible
;       OPERATING SYSTEM: PC-DOS/MS-DOS
;       SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)
;
;   CURRENT APPLICATION DATA:
;       APPLICATION NUMBER: US/09/938,703
;       FILING DATE: 24-Aug-2001
;       CLASSIFICATION: <Unknown>
;
;   PRIOR APPLICATION DATA:
;       APPLICATION NUMBER: 09/626,939
;       FILING DATE: 2000-07-27
;
;   ATTORNEY/AGENT INFORMATION:
;       NAME: Altman, Daniel E
;       REGISTRATION NUMBER: 34,115
;       REFERENCE/DOCKET NUMBER: <Unknown>
;
;   INFORMATION FOR SEQ ID NO: 5:
;       SEQUENCE CHARACTERISTICS:
;           LENGTH: 352 amino acids
;           TYPE: amino acid
;           TOPOLOGY: linear
;       MOLECULE TYPE: protein
;       SEQUENCE DESCRIPTION: SEQ ID NO: 5:
US-09-938-703-5

```

```

Query Match          42.9%;  Score 379;  DB 9;  Length 352;
Best Local Similarity 35.0%;  Pred. No. 1.8e-29;
Matches 99;  Conservative 7;  Mismatches 27;  Indels 150;  Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          | | | ||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||| : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

Qy      125 -----GGSYKCGLC----QEFG LN NCSSSNRLDGHQRV 154
          | | ||| : |
Db      239 TIMIVYFLFWAPYNIVLLNTFQEFG LN NCSSSNRLDQAMQV 281

```

RESULT 15
 US-09-502-783A-2
 ; Sequence 2, Application US/09502783A
 ; Patent No. US20020132269A1
 ; GENERAL INFORMATION:
 ; APPLICANT: Li, Yi
 ; APPLICANT: Ruben, Steven M.
 ; TITLE OF INVENTION: Polynucleotides Encoding Human G-Protein Chemokine
 Receptor (CCR5)
 ; TITLE OF INVENTION: HDG NR10
 ; FILE REFERENCE: 1488.1150006
 ; CURRENT APPLICATION NUMBER: US/09/502,783A
 ; CURRENT FILING DATE: 2001-08-23
 ; PRIOR APPLICATION NUMBER: 08/466,343
 ; PRIOR FILING DATE: 1995-06-06
 ; NUMBER OF SEQ ID NOS: 9
 ; SOFTWARE: PatentIn version 3.0
 ; SEQ ID NO 2
 ; LENGTH: 352
 ; TYPE: PRT
 ; ORGANISM: Homo sapiens
 US-09-502-783A-2

Query Match 42.9%; Score 379; DB 9; Length 352;
 Best Local Similarity 35.0%; Pred. No. 1.8e-29;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||
Db      1 MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |  |                      ||| : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||
Db      119 IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||| : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |  | ||| :|
Db      239 TIMIVYFLEWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

Search completed: March 5, 2004, 16:33:45
 Job time : 85.7994 secs

GenCore version 5.1.6
Copyright (c) 1993 - 2004 Compugen Ltd.

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:15:44 ; Search time 104.182 Seconds
(without alignments)
475.479 Million cell updates/sec

Title: US-10-057-890A-31
Perfect score: 884
Sequence: 1 MKVSVAAALSCMLMLVTALGSM.....GLNNCSSSNRLDGHQRVHAA 157

Scoring table: BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 1017041 seqs, 315518202 residues

Total number of hits satisfying chosen parameters: 1017041

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : SPTREMBL_25:*
1: sp_archaea:*
2: sp_bacteria:*
3: sp_fungi:*
4: sp_human:*
5: sp_invertebrate:*
6: sp_mammal:*
7: sp_mhc:*
8: sp_organelle:*
9: sp_phage:*
10: sp_plant:*
11: sp_rodent:*
12: sp_virus:*
13: sp_vertebrate:*
14: sp_unclassified:*
15: sp_rvirus:*
16: sp_bacteriap:*
17: sp_archeap:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result	Score	Query	Match	Length	DB	ID	Description

1	373	42.2	352	6	O18772	O18772	pan troglod
2	370	41.9	344	6	Q9TQR8	Q9tqr8	cercocebus
3	370	41.9	344	6	O77833	O77833	cercocebus
4	369	41.7	352	6	Q95NC5	Q95nc5	hylobates s
5	369	41.7	352	6	Q9TV50	Q9tv50	pan troglod
6	367	41.5	352	6	Q95NC3	Q95nc3	miopithecus
7	367	41.5	352	6	O18771	O18771	pan troglod
8	366	41.4	352	6	Q9TSK1	Q9tsk1	cercopithec
9	366	41.4	352	6	Q9TV49	Q9tv49	cercocebus
10	366	41.4	352	6	Q9XT13	Q9xt13	papio anubi
11	366	41.4	352	6	Q9TV45	Q9tv45	cercopithec
12	366	41.4	352	6	Q95NE8	Q95ne8	cercopithec
13	365	41.3	352	6	Q9XT14	Q9xt14	colobus gue
14	365	41.3	352	6	Q95NC6	Q95nc6	trachypithe
15	365	41.3	352	6	Q95NC8	Q95nc8	colobus pol
16	364	41.2	352	6	Q9XS99	Q9xs99	gorilla gor
17	363	41.1	352	6	O97975	O97975	macaca arct
18	363	41.1	352	6	Q9MZA2	Q9mza2	cercopithec
19	362	41.0	352	6	Q95NC1	Q95nc1	theropithec
20	362	41.0	352	6	Q95NC0	Q95nc0	hylobates m
21	362	41.0	352	6	Q9TV46	Q9tv46	cercopithec
22	362	41.0	352	6	O97962	O97962	pygathrix a
23	361	40.8	352	6	Q95NC7	Q95nc7	nasalis lar
24	361	40.8	352	6	Q9TV42	Q9tv42	cercopithec
25	361	40.8	352	6	O77776	O77776	cercocebus
26	361	40.8	352	6	Q95NE1	Q95ne1	cercocebus
27	361	40.8	352	6	Q9TQX0	Q9tqx0	cercopithec
28	359	40.6	352	6	Q9TV43	Q9tv43	cercopithec
29	358	40.5	352	6	O18770	O18770	pan troglod
30	357	40.4	352	6	Q9TV47	Q9tv47	cercopithec
31	357	40.4	352	6	Q9BGN5	Q9bgn5	cercopithec
32	357	40.4	352	6	Q9XT12	Q9xt12	cercopithec
33	357	40.4	352	6	Q9TSQ7	Q9tsq7	cercopithec
34	356	40.3	352	6	Q95ND2	Q95nd2	mandrillus
35	356	40.3	352	6	Q95ND1	Q95nd1	mandrillus
36	355	40.2	352	6	Q9XT76	Q9xt76	cercopithec
37	355	40.2	352	6	Q9TV44	Q9tv44	cercopithec
38	355	40.2	352	6	Q9XS35	Q9xs35	macaca neme
39	355	40.2	352	6	Q9MZA3	Q9mza3	hylobates a
40	355	40.2	352	6	Q9TV93	Q9tv93	macaca arct
41	355	40.2	352	6	Q95ND0	Q95nd0	erythrocebu
42	346	39.1	352	6	Q9BGN6	Q9bgn6	cercopithec
43	345	39.0	339	4	Q9UN25	Q9un25	homo sapien
44	344	38.9	339	4	Q9UN23	Q9un23	homo sapien
45	344	38.9	339	4	Q9UN27	Q9un27	homo sapien

ALIGNMENTS

RESULT 1

O18772

ID O18772 PRELIMINARY; PRT; 352 AA.

AC O18772;

DT 01-JAN-1998 (TrEMBLrel. 05, Created)

DT 01-JAN-1998 (TrEMBLrel. 05, Last sequence update)

DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)

DE CCR5 receptor (Fragment).
 GN CCR5.
 OS Pan troglodytes (Chimpanzee).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Pan.
 OX NCBI_TaxID=9598;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC STRAIN=ChCCR5-142a;
 RA Zhang L., Carruthers C.D., He T., Huang Y., Cao Y., Wang G., Hahn B.,
 RA Ho D.D.;
 RT "HIV-1 subtypes, co-receptor usage, and CCR5 polymorphism.";
 RL AIDS Res. Hum. Retroviruses 0:0-0(1997).
 DR EMBL; AF011541; AAB65741.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 FT NON TER 352 352
 SQ SEQUENCE 352 AA; 40598 MW; 39688FA7004C952F CRC64;

Query Match 42.2%; Score 373; DB 6; Length 352;
 Best Local Similarity 34.6%; Pred. No. 1.7e-33;
 Matches 98; Conservative 8; Mismatches 27; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 |||||:|||||
 Db 1 MDYQVSSPIYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVDNMLVILILINCKR 60

 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVTVSVITWVVAVFASLPGIIFTRSQKEGLHYTC 178

 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRAVRLIF 238

 Qy 125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYDIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 2

Q9TQR8

ID Q9TQR8 PRELIMINARY; PRT; 344 AA.

AC Q9TQR8;

DT 01-MAY-2000 (TrEMBLrel. 13, Created)

DT 01-MAY-2000 (TrEMBLrel. 13, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE Chemokine receptor CCR5.
 GN CCR5.
 OS Cercocebus torquatus torquatus, and
 OS Cercocebus torquatus atys (Red-crowned mangabey) (Sooty mangabey).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Cercocebus.
 OX NCBI_TaxID=81944, 9531;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC SPECIES=C.torquatus torquatus; STRAIN=1049, and 997;
 RA Chen Z., Kwon D., Jin Z., Monard S., Telfer P., Jones M., Lu C.,
 RA Aguilar R., Ho D.D., Marx P.A.;
 RT "Natural infection of a homozygous delta 24 CCR5 red-capped mangabey
 RT with a R2b-tropic simian immunodeficiency virus.";
 RL J. Exp. Med. 0:0-0(1998).
 RN [2]
 RP SEQUENCE FROM N.A.
 RC SPECIES=C.torquatus atys;
 RA Palacios E., Digilio L., McClure H.M., Chen Z., Marx P.A.,
 RA Goldsmith M.A., Grant R.M.;
 RT "Parallel evolution of CCR5-null phenotypes in humans and in a natural
 RT host of simian immunodeficiency virus.";
 RL Curr. Biol. 0:0-0(1998).
 RN [3]
 RP SEQUENCE FROM N.A.
 RC SPECIES=C.torquatus torquatus; STRAIN=RCM411;
 RA Beer B.E., Kuiken C.L., Tooze Z., Foley B.T., Goeken R.M., Brown C.R.,
 RA St Claire M., Hirsch V.M.;
 RT "Characterization of novel simian immunodeficiency viruses from
 RT redcapped mangabeys from Nigeria (SIVrcmNgM411 and NgD409).";
 RL Submitted (FEB-2001) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF094753; AAC62474.1; -.
 DR EMBL; AF079473; AAC31194.1; -.
 DR EMBL; AF084003; AAC62471.1; -.
 DR EMBL; AF349683; AAK69685.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 344 AA; 39592 MW; E15F5F601191A4D1 CRC64;

Query Match 41.9%; Score 370; DB 6; Length 344;
 Best Local Similarity 34.9%; Pred. No. 3.5e-33;
 Matches 96; Conservative 9; Mismatches 28; Indels 142; Gaps 7;

Qy 20 MDYQVSSPIYDINYTTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||||
 Db 1 MDYQVSSPTYDIDYTTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60

```

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
Db      61 LKSMTDIYLLNLAIISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTCSSHFYPYSQ 105
      : | | | | | | | | | | | | | | | | | | | | | | | | | | | |
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVAVFASLPGIIFTRSQREGLHYTCSSHFYPYSQ 178

Qy     106 YQFWKNFQTLKI-----HQRVHGG----- 124
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
Db     179 YQFWKNFQTLKIVILGLVPLLVVICYSGILKTLRLCRNEKKRHRVRLIFTIMIVYFL 238

Qy     125 -GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
Db     239 FWAPYNIVLLLNTFQEFGFLNNCSSSNRLDQAMQV 273

```

RESULT 3
O77833

```

ID   O77833          PRELIMINARY;          PRT;    344 AA.
AC   O77833;
DT   01-NOV-1998 (TrEMBLrel. 08, Created)
DT   01-NOV-1998 (TrEMBLrel. 08, Last sequence update)
DT   01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
DE   Chemokine receptor CCR5.
GN   CCR5.
OS   Cercopithecus torquatus torquatus.
OC   Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC   Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
OC   Cercopithecinae; Cercopithecus.
OX   NCBI_TaxID=81944;
RN   [1]
RP   SEQUENCE FROM N.A.
RC   STRAIN=1208, and 009;
RA   Chen Z., Kwon D., Jin Z., Monard S., Telfer P., Jones M., Lu C.,
RA   Aguilar R., Ho D.D., Marx P.A.;
RT   "Natural infection of a homozygous delta 24 CCR5 red-capped mangabey
RT   with a R2b-tropic simian immunodeficiency virus.";
RL   J. Exp. Med. 0:0-0(1998).
DR   EMBL; AF094752; AAC62473.1; -.
DR   EMBL; AF084002; AAC62470.1; -.
DR   GO; GO:0016021; C:integral to membrane; IEA.
DR   GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
DR   GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
DR   InterPro; IPR000276; GPCR_Rhodpsn.
DR   Pfam; PF00001; 7tm_1; 1.
DR   PRINTS; PR00237; GPCRRHODOPSN.
DR   PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR   PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
SQ   SEQUENCE   344 AA;  39578 MW;  0CEC05E47C2F6DE6 CRC64;

```

```

Query Match          41.9%;  Score 370;  DB 6;  Length 344;
Best Local Similarity 34.9%;  Pred. No. 3.5e-33;
Matches 96;  Conservative 9;  Mismatches 28;  Indels 142;  Gaps 7;

```

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49

```



```

      ||||| |||:|||||
Db      1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFVFGFVGNILVVLILINCKR 60
Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | ||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
Qy      84 -----LCTRSQKEGLHYTCSSHPYQS 105
      : ||||:|||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVAVFASLPGIIFTRSQREGLHYTCSSHPYQS 178
Qy      106 YQFWKNFQTLKI-----HQRVHGG----- 124
      ||||| ||| : |
Db      179 YQFWKNFQTLKIVILGLVPLLMVICYSGILKTLRLCRNEKKRHRAVRLIFTIMIVYFL 238
Qy      125 -GGSYKCGLC----QEFGNLCSSSNRLDGHQRV 154
      | | ||||| ||| : |
Db      239 FWAPYNIVLLNTFQEFGNLCSSSNRLDQAMQV 273

```

RESULT 4

Q95NC5

```

ID   Q95NC5          PRELIMINARY;      PRT;   352 AA.
AC   Q95NC5;
DT   01-DEC-2001 (TrEMBLrel. 19, Created)
DT   01-DEC-2001 (TrEMBLrel. 19, Last sequence update)
DT   01-OCT-2003 (TrEMBLrel. 25, Last annotation update)
DE   C-C chemokine receptor 5.
GN   CCR5.
OS   Hylobates syndactylus (Siamang) (Symphalangus syndactylus).
OC   Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC   Mammalia; Eutheria; Primates; Catarrhini; Hylobatidae; Hylobates.
OX   NCBI_TaxID=9590;
RN   [1]
RP   SEQUENCE FROM N.A.
RA   Zhang Y., Ryder O.A., Zhang Y.;
RT   "Sequence comparison of the CCR5 gene in primates and primate
RT   phylogeny.";
RL   Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
DR   EMBL; AF177884; AAK43367.1; -.
DR   GO; GO:0016021; C:integral to membrane; IEA.
DR   GO; GO:0004872; F:receptor activity; IEA.
DR   GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
DR   GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
DR   InterPro; IPR000276; GPCR_Rhodpsn.
DR   Pfam; PF00001; 7tm_1; 1.
DR   PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR   PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
KW   Receptor.
SQ   SEQUENCE   352 AA;  40508 MW;  F4F64B3AD5AF658A CRC64;

```

Query Match 41.7%; Score 369; DB 6; Length 352;

Best Local Similarity 34.3%; Pred. No. 4.7e-33;

Matches 97; Conservative 8; Mismatches 28; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      ||||| |||:|||||

```

```

Db          1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
QY          50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHSHHSYKCG--- 83
              |   |               ||||| | | | | | | | | | | | : | :   : |
Db          61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
QY          84 -----LCTRSQKEGLHYTC 97
              : ||||| | | | | | | | | | | |
Db          119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178
QY          98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
              ||||| | | | | | | | | | | | : : |
Db          179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRAVRLIF 238
QY          125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
              |   |               ||||| | | | | | | | | | | | : |
Db          239 TIMIVYFLFWAPYNIVLLNLTFFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 5

Q9TV50

```

ID   Q9TV50          PRELIMINARY;          PRT;    352 AA.
AC   Q9TV50;
DT   01-MAY-2000 (TrEMBLrel. 13, Created)
DT   01-MAY-2000 (TrEMBLrel. 13, Last sequence update)
DT   01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
DE   CC chemokine receptor type 5.
GN   CCR5.
OS   Pan troglodytes (Chimpanzee).
OC   Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC   Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Pan.
OX   NCBI_TaxID=9598;
RN   [1]
RP   SEQUENCE FROM N.A.
RC   STRAIN=1410;
RX   MEDLINE=99335215; PubMed=10408730;
RA   Muller-Trutwin M.C., Corbet S., Hansen J., Georges-Courbot M.C.,
RA   Diop O., Rigoulet J., Barre-Sinoussi F., Fomsgaard A.;
RT   "Mutations in CCR5-coding sequences are not associated with SIV
RT   carrier status in African nonhuman primates.";
RL   AIDS Res. Hum. Retroviruses 15:931-939(1999).
RN   [2]
RP   SEQUENCE FROM N.A.
RC   STRAIN=1410;
RA   Mueller-Trutwin M.C., Corbet S., Hansen J., Diop O.,
RA   Georges-Courbot M.-C., Barre-Sinoussi F., Fomsgaard A.;
RL   Submitted (NOV-1997) to the EMBL/GenBank/DBJ databases.
DR   EMBL; AF035214; AAD44007.1; -.
DR   GO; GO:0016021; C:integral to membrane; IEA.
DR   GO; GO:0004872; F:receptor activity; IEA.
DR   GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
DR   GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
DR   InterPro; IPR000276; GPCR_Rhodpsn.
DR   Pfam; PF00001; 7tm_1; 1.
DR   PRINTS; PR00237; GPCRRHODOPSN.
DR   PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR   PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.

```

SQ SEQUENCE 352 AA; 40481 MW; 2578A0F2C07A4A65 CRC64;

Query Match 41.7%; Score 369; DB 6; Length 352;
Best Local Similarity 34.3%; Pred. No. 4.7e-33;
Matches 97; Conservative 8; Mismatches 28; Indels 150; Gaps 7;

Qy	20	MDYQVSSPIYDINYTSEPCQKINVKQIAA-----	49
		:	
Db	1	MDYQVSSPTYDIDYTTSEPCQKINVKQIAARLLPPLYSLVFI FGFVGNILVVLILINCKR	60
Qy	50	-----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG---	83
		: :	
Db	61	LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF	118
Qy	84	-----LCTRSQKEGLHYTC	97
		:	
Db	119	IILLSIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQKEGLHYTC	178
Qy	98	SSHFPYSQYQFWKNFQTLKI-----HQRVHGG-----	124
		: :	
Db	179	SSHFPYSQYQFWKNFQTLKIVILGLVLPLLVMVICYSGILKTLLRCRNEKKRHRVRLIF	238
Qy	125	-----GGSYKCGLC----QEFGNLCSSSNRLDGHQRV	154
		:	
Db	239	TIMIVYFLFWAPYNIVLLNTFQEFFGLNLCSSSNRLDOAMOV	281

095NC3

```

ID      Q95NC3          PRELIMINARY;          PRT;    352 AA.
AC      Q95NC3;
DT      01-DEC-2001 (TrEMBLrel. 19, Created)
DT      01-DEC-2001 (TrEMBLrel. 19, Last sequence update)
DT      01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
DE      C-C chemokine receptor 5.
GN      CCR5.
OS      Miopithecus talapoin (Talapoin) (Cercopithecus talapoin).
OC      Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC      Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
OC      Cercopithecinae; Miopithecus.
OX      NCBI_TaxID=36231;
RN      [1]
RP      SEQUENCE FROM N.A.
RA      Zhang Y., Ryder O.A., Zhang Y.;
RT      "Sequence comparison of the CCR5 gene in primates and primate
RT      phylogeny.";
RL      Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
DR      EMBL; AF177886; AAK43369.1; -.
DR      GO; GO:0016021; C:integral to membrane; IEA.
DR      GO; GO:0004872; F:receptor activity; IEA.
DR      GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
DR      GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
DR      InterPro; IPR000276; GPCR_Rhodpsn.
DR      Pfam; PF00001; 7tm_1; 1.
DR      PRINTS; PR00237; GPCRRHODOPSN.
DR      PROSITE; PS00237; G PROTEIN RECEPTOR F1 1; 1.

```

DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40546 MW; 6464152F3E566AE5 CRC64;

Query Match 41.5%; Score 367; DB 6; Length 352;
 Best Local Similarity 33.9%; Pred. No. 7.8e-33;
 Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

```

QY      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||| ||||| ||||| ||||| ||||| |||||
Db      1 MDYQVSSPTYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60

QY      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
          |   |               ||||| ||||| : : | :   |
Db      61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--RLLTGLYFIGFFSGIFF 118

QY      84 -----LCTRSQKEGLHYTC 97
          : |||: |||||
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQREGLHYTC 178

QY      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||| ||||| ||||| ||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRVRLIF 238

QY     125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
          |   |   ||||| ||||| ||||| : |
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

RESULT 7

O18771

ID O18771 PRELIMINARY; PRT; 352 AA.
 AC O18771;
 DT 01-JAN-1998 (TrEMBLrel. 05, Created)
 DT 01-JAN-1998 (TrEMBLrel. 05, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE CCR5 receptor (Fragment).
 GN CCR5.
 OS Pan troglodytes (Chimpanzee).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Pan.
 OX NCBI_TaxID=9598;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC STRAIN=ChCCR5-141a;
 RA Zhang L., Carruthers C.D., He T., Huang Y., Cao Y., Wang G., Hahn B.,
 RA Ho D.D.;
 RT "HIV-1 subtypes, co-receptor usage, and CCR5 polymorphism.";
 RL AIDS Res. Hum. Retroviruses 0:0-0(1997).
 DR EMBL; AF011539; AAB65739.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.

DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 FT NON_TER 352 352
 SQ SEQUENCE 352 AA; 40466 MW; 3FFFAC7ABAE1D4FB CRC64;

Query Match 41.5%; Score 367; DB 6; Length 352;
 Best Local Similarity 34.3%; Pred. No. 7.8e-33;
 Matches 97; Conservative 8; Mismatches 28; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||:|||||
Db      1 MDYQVSSPIYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          | | ||||| | : | : |
Db      61 LKSMTDIYLLNLAINLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||||
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKGGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||| |||||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVPLLVVICYSGILKTLRLCRNEKKRHRAVRLIF 238

Qy     125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
          | | ||||| :|
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

RESULT 8

Q9TSK1

ID Q9TSK1 PRELIMINARY; PRT; 352 AA.
 AC Q9TSK1;
 DT 01-MAY-2000 (TrEMBLrel. 13, Created)
 DT 01-MAY-2000 (TrEMBLrel. 13, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE G-protein coupled chemokine receptor.
 OS Cercopithecus aethiops (Green monkey) (Grivet).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Cercopithecus.
 OX NCBI_TaxID=9534;
 RN [1]
 RP SEQUENCE FROM N.A.
 RA Holtkamp N., Baier M., Werner A.;
 RL Submitted (AUG-1997) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF019379; AAD01639.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.

DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
KW Receptor.
SQ SEQUENCE 352 AA; 40588 MW; 0F1869D9A6668DBB CRC64;

Query Match 41.4%; Score 366; DB 6; Length 352;
Best Local Similarity 33.9%; Pred. No. 1e-32;
Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

```
Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          ||||| |||:|||||
Db      1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFI GFVGNILVVLILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      ||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||:|||||
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPRIIFTRSQREGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          ||||| |||||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVPLLVVICYSGLKTLRLCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC-----QEFFGLNNCSSSNRLDGHQRV 154
          |  |                      ||||| :|
Db     239 TIMIVYFLEWAPYNIVLLNLTQEFFGLNNCSSSNRLDQAMQV 281
```

RESULT 9

Q9TV49

ID Q9TV49 PRELIMINARY; PRT; 352 AA.
AC Q9TV49;
DT 01-MAY-2000 (TrEMBLrel. 13, Created)
DT 01-MAY-2000 (TrEMBLrel. 13, Last sequence update)
DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
DE CC chemokine receptor type 5 (C-C chemokine receptor 5).
GN CCR5.
OS Cercopithecus galeries (Agile mangabey).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
OC Cercopithecinae; Cercopithecus.
OX NCBI_TaxID=9532;
RN [1]
RP SEQUENCE FROM N.A.
RC STRAIN=4;
RX MEDLINE=99335215; PubMed=10408730;
RA Muller-Trutwin M.C., Corbet S., Hansen J., Georges-Courbot M.C.,
RA Diop O., Rigoulet J., Barre-Sinoussi F., Fomsgaard A.;
RT "Mutations in CCR5-coding sequences are not associated with SIV
RT carrier status in African nonhuman primates."
RL AIDS Res. Hum. Retroviruses 15:931-939(1999).
RN [2]
RP SEQUENCE FROM N.A.
RC STRAIN=4;
RA Mueller-Trutwin M.C., Corbet S., Hansen J., Diop O.,
RA Georges-Courbot M.-C., Barre-Sinoussi F., Fomsgaard A.;

RL Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
 RN [3]
 RP SEQUENCE FROM N.A.
 RA Zhang Y., Ryder O.A., Zhang Y.;
 RT "Sequence comparison of the CCR5 gene in primates and primate
 RT phylogeny."
 RL Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF035215; AAD44008.1; -.
 DR EMBL; AF177898; AAK43381.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40479 MW; 5E1504A9B87278B2 CRC64;

Query Match 41.4%; Score 366; DB 6; Length 352;
 Best Local Similarity 33.9%; Pred. No. 1e-32;
 Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLLFLLTVPFWAHAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGLIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| ||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVPLLVVICYSGLKTLRLCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC---QEFGFLNNCSSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLLNTFQEFGFLNNCSSSNRLDQAMQV 281

RESULT 10

Q9XT13

ID Q9XT13 PRELIMINARY; PRT; 352 AA.
 AC Q9XT13;
 DT 01-NOV-1999 (TrEMBLrel. 12, Created)
 DT 01-NOV-1999 (TrEMBLrel. 12, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE CC chemokine receptor 5.
 GN CCR5.
 OS Papio anubis (Olive baboon).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Papio.
 OX NCBI_TaxID=9555;
 RN [1]
 RP SEQUENCE FROM N.A.
 RA Spearman P.W., Mburu D.N., Graham B.S.;
 RT "Differential Utilization of CCR5 Molecules from Three East African
 RT Simian Species by the HIV-1 Envelope Glycoprotein."
 RL Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF141640; AAD32685.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40489 MW; 0B47E337C11E2E1E CRC64;

Query Match 41.4%; Score 366; DB 6; Length 352;
 Best Local Similarity 33.9%; Pred. No. 1e-32;
 Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDNYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : ||||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| ||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVPLLVVICYSGLKTLRCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLNLTFFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 11

Q9TV45

ID Q9TV45 PRELIMINARY; PRT; 352 AA.
 AC Q9TV45;
 DT 01-MAY-2000 (TrEMBLrel. 13, Created)
 DT 01-MAY-2000 (TrEMBLrel. 13, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE CC chemokine receptor type 5.
 GN CCR5.
 OS Cercopithecus nictitans (white-nosed guenon).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Cercopithecus.
 OX NCBI_TaxID=36228;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC STRAIN=7;
 RX MEDLINE=99335215; PubMed=10408730;
 RA Muller-Trutwin M.C., Corbet S., Hansen J., Georges-Courbot M.C.,
 RA Diop O., Rigoulet J., Barre-Sinoussi F., Fomsgaard A.;
 RT "Mutations in CCR5-coding sequences are not associated with SIV
 RT carrier status in African nonhuman primates.";
 RL AIDS Res. Hum. Retroviruses 15:931-939(1999).
 RN [2]
 RP SEQUENCE FROM N.A.
 RC STRAIN=7;
 RA Mueller-Trutwin M.C., Corbet S., Hansen J., Diop O.,
 RA Georges-Courbot M.-C., Barre-Sinoussi F., Fomsgaard A.;
 RL Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF035219; AAD44012.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40658 MW; BC665891ED5661F8 CRC64;

Query Match 41.4%; Score 366; DB 6; Length 352;

Best Local Similarity 33.9%; Pred. No. 1e-32;

Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
          |||||  |||:|||||  |||||
Db      1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPLLYSLVFIFGFVGNILVVLILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
          |  |                      |||||  |||  | : | : |
Db      61 LKSMTDIYLLNLAIISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
          : |||:|||||
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPRIIFTRSQREGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
          |||||  |||||  |||||  |||||  |||||  |||||  |||||  |||||  |||||  |||||
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLPVLLVMVICYSGLKTLRLCRNEKKRHRVAMLIF 238

Qy     125 -----GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
          |  |                      |||||  |||||  |||||  |||||  |||||  |||||
Db     239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281
  
```

RESULT 12

Q95NE8

ID Q95NE8 PRELIMINARY; PRT; 352 AA.
 AC Q95NE8;
 DT 01-DEC-2001 (TrEMBLrel. 19, Created)
 DT 01-DEC-2001 (TrEMBLrel. 19, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE CC chemokine receptor type 5.
 GN CCR5.
 OS Cercopithecus tantalus.
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Cercopithecus.
 OX NCBI_TaxID=60712;
 RN [1]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=99335215; PubMed=10408730;
 RA Muller-Trutwin M.C., Corbet S., Hansen J., Georges-Courbot M.C.,
 RA Diop O., Rigoulet J., Barre-Sinoussi F., Fomsgaard A.;
 RT "Mutations in CCR5-coding sequences are not associated with SIV
 RT carrier status in African nonhuman primates.";
 RL AIDS Res. Hum. Retroviruses 15:931-939(1999).
 DR EMBL; AF081577; AAD45495.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 SQ SEQUENCE 352 AA; 40576 MW; 0F1869D9B7968DBB CRC64;

Query Match 41.4%; Score 366; DB 6; Length 352;

Best Local Similarity 33.9%; Pred. No. 1e-32;

Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVQIAA----- 49
 ||||| |||:|||||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : ||||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPRIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| ||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLLRCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 13

Q9XT14

ID Q9XT14 PRELIMINARY; PRT; 352 AA.
 AC Q9XT14;
 DT 01-NOV-1999 (TrEMBLrel. 12, Created)
 DT 01-NOV-1999 (TrEMBLrel. 12, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE CC chemokine receptor 5.
 GN CCR5.
 OS Colobus guereza (Black-and-white colobus monkey).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;
 OC Colobus.
 OX NCBI_TaxID=33548;
 RN [1]
 RP SEQUENCE FROM N.A.
 RA Spearman P.W., Mburu D.N., Graham B.S.;
 RT "Differential Utilization of CCR5 Molecules from Three East African
 RT Simian Species by the HIV-1 Envelope Glycoprotein."
 RL Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF141639; AAD32684.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40550 MW; 9B078EF04D34DB36 CRC64;

Query Match 41.3%; Score 365; DB 6; Length 352;

Best Local Similarity 33.6%; Pred. No. 1.3e-32;

Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVQIAA----- 49
 ||||| :|||:|||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKVNQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTATFGVVTSVITWVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRAVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLNTFQEFFGLNNC SSSNRLDQAMQV 281

RESULT 14

Q95NC6

ID Q95NC6 PRELIMINARY; PRT; 352 AA.
 AC Q95NC6;
 DT 01-DEC-2001 (TrEMBLrel. 19, Created)
 DT 01-DEC-2001 (TrEMBLrel. 19, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE C-C chemokine receptor 5.
 GN CCR5.
 OS Trachypithecus johnii (hooded leaf monkey).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;
 OC Trachypithecus.
 OX NCBI_TaxID=66063;
 RN [1]
 RP SEQUENCE FROM N.A.
 RA Zhang Y., Ryder O.A., Zhang Y.;
 RT "Sequence comparison of the CCR5 gene in primates and primate
 RT phylogeny."
 RL Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF177883; AAK43366.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40462 MW; 52824E0322559F7F CRC64;

Query Match 41.3%; Score 365; DB 6; Length 352;

Best Local Similarity 33.6%; Pred. No. 1.3e-32;

Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYTSEPCQKINVQIAA----- 49
 ||||| :|||:|||||
 Db 1 MDYQVSSPTYDIDYTTSEPCQKVNKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
 | | ||||| : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLPPLVMVICYSGLKTLRLCRSEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
 | | ||||| : |

RESULT 15

Q95NC8

ID Q95NC8 PRELIMINARY; PRT; 352 AA.
 AC Q95NC8;
 DT 01-DEC-2001 (TrEMBLrel. 19, Created)
 DT 01-DEC-2001 (TrEMBLrel. 19, Last sequence update)
 DT 01-JUN-2003 (TrEMBLrel. 24, Last annotation update)
 DE C-C chemokine receptor 5.
 GN CCR5.
 OS Colobus polykomos.
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;
 OC Colobus.
 OX NCBI_TaxID=9572;
 RN [1]
 RP SEQUENCE FROM N.A.
 RA Zhang Y., Ryder O.A., Zhang Y.;
 RT "Sequence comparison of the CCR5 gene in primates and primate
 RT phylogeny."
 RL Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
 DR EMBL; AF177881; AAK43364.1; -.
 DR GO; GO:0016021; C:integral to membrane; IEA.
 DR GO; GO:0004872; F:receptor activity; IEA.
 DR GO; GO:0001584; F:rhodopsin-like receptor activity; IEA.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; IEA.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW Receptor.
 SQ SEQUENCE 352 AA; 40578 MW; 4366F149C3B4938F CRC64;

Query Match 41.3%; Score 365; DB 6; Length 352;

Best Local Similarity 33.6%; Pred. No. 1.3e-32;

Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||:|||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKVNKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| |||||
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVPLLVVICYSGLKTLRLCRNEKKRHRAVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154

Search completed: March 5, 2004, 16:27:34
Job time : 106.182 secs

OM protein - protein search, using sw model

Run on: March 5, 2004, 16:15:14 ; Search time 21.321 Seconds
(without alignments)
383.426 Million cell updates/sec

Title: US-10-057-890A-31
Perfect score: 884
Sequence: 1 MKVSVAALSCLMLVTALGSM.....GLNNCSSSRLDGHQVRHAA 157

Scoring table: BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 141681 seqs, 52070155 residues

Total number of hits satisfying chosen parameters: 141681

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : SwissProt_42:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Query		DB	ID	Description
	Score	Match Length			
1	379	42.9 352	1	CKR5_HUMAN	P51681 homo sapien
2	374	42.3 352	1	CKR5_PANTR	P56440 pan troglod
3	369	41.7 352	1	CKR5_HYLSY	Q95nc5 hylobates s
4	369	41.7 352	1	CKR5_PONPY	O97881 pongo pygma
5	366	41.4 352	1	CKR5_PAPHA	P56441 papio hamad
6	365	41.3 352	1	CKR5_PYGBI	O97880 pygathrix b
7	365	41.3 352	1	CKR5_PYGNE	O97882 pygathrix n
8	365	41.3 352	1	CKR5_TRAFR	O97878 trachypithe
9	365	41.3 352	1	CKR5_TRAPH	O97879 trachypithe
10	364	41.2 352	1	CKR5_GORG	P56439 gorilla gor
11	363	41.1 352	1	CKR5_MACMU	P79436 macaca mula
12	362	41.0 352	1	CKR5_HYLM	Q95nc0 hylobates m
13	361	40.8 352	1	CKR5_CERP	Q9tv42 cercopithec
14	361	40.8 352	1	CKR5_CERTO	O62743 cercocobus
15	359	40.6 352	1	CKR5_HYLL	O97883 hylobates l
16	357	40.4 352	1	CKR5_CERAE	P56493 cercopithec
17	272	30.8 354	1	CKR5_MOUSE	P51682 mus musculu

18	249	28.2	354	1	CKR5_RAT	008556	rattus norv
19	156.5	17.7	474	1	Z256_HUMAN	Q9y2p7	homo sapien
20	154	17.4	201	1	Z239_MOUSE	P24399	mus musculu
21	151.5	17.1	645	1	Z235_MOUSE	Q61116	mus musculu
22	151.5	17.1	754	1	Z287_HUMAN	Q9hbt7	homo sapien
23	150.5	17.0	501	1	ZF96_MOUSE	Q9zld7	mus musculu
24	146.5	16.6	458	1	Z239_HUMAN	Q16600	homo sapien
25	144.5	16.3	759	1	Z287_MOUSE	Q9eqb9	mus musculu
26	144	16.3	698	1	Z234_HUMAN	Q14588	homo sapien
27	140.5	15.9	604	1	Z305_HUMAN	O43309	homo sapien
28	140.5	15.9	744	1	YJ62_HUMAN	Q8tf39	homo sapien
29	140.5	15.9	803	1	Z226_HUMAN	Q9nyt6	homo sapien
30	140.5	15.9	913	1	Z228_HUMAN	Q9uju3	homo sapien
31	140	15.8	578	1	Z192_HUMAN	Q15776	homo sapien
32	139.5	15.8	682	1	ZN45_HUMAN	Q02386	homo sapien
33	139.5	15.8	751	1	Z184_HUMAN	Q99676	homo sapien
34	139	15.7	810	1	Z33A_HUMAN	Q06730	homo sapien
35	138.5	15.7	468	1	ZF90_HUMAN	Q8tf47	homo sapien
36	138.5	15.7	670	1	ZN16_HUMAN	P17020	homo sapien
37	138.5	15.7	1029	1	Z197_HUMAN	O14709	homo sapien
38	138	15.6	463	1	Z331_HUMAN	Q9nqx6	homo sapien
39	136.5	15.4	348	1	Z134_HUMAN	P52741	homo sapien
40	136.5	15.4	659	1	Z304_HUMAN	Q9hcx3	homo sapien
41	136.5	15.4	825	1	ZF28_MOUSE	P10078	mus musculu
42	135.5	15.3	643	1	ZN74_HUMAN	Q16587	homo sapien
43	133.5	15.1	636	1	ZF90_MOUSE	Q61967	mus musculu
44	132.5	15.0	642	1	ZN14_HUMAN	P17017	homo sapien
45	131.5	14.9	367	1	Z211_HUMAN	Q13398	homo sapien

ALIGNMENTS

RESULT 1

CKR5_HUMAN

ID CKR5_HUMAN STANDARD; PRT; 352 AA.

AC P51681; O14692; O14693; O14695; O14696; O14697; O14698; O14699;

AC O14700; O14701; O14702; O14703; O14704; O14705; O14706; O14707;

AC O14708; O15538; Q9UPA4;

DT 01-OCT-1996 (Rel. 34, Created)

DT 01-OCT-1996 (Rel. 34, Last sequence update)

DT 15-MAR-2004 (Rel. 43, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5)

DE (HIV-1 fusion co-receptor) (CHEMR13) (CD195 antigen).

GN CCR5 OR CMKBR5.

OS Homo sapiens (Human).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

OX NCBI_TaxID=9606;

RN [1]

RP SEQUENCE FROM N.A.

RX MEDLINE=96241590; PubMed=8639485;

RA Samson M., Labbe O., Mollereau C., Vassart G., Parmentier M.;

RT "Molecular cloning and functional expression of a new human

RT CC-chemokine receptor gene.";

RL Biochemistry 35:3362-3367(1996).

RN [2]

RP SEQUENCE FROM N.A.
 RX MEDLINE=96291862; PubMed=8663314;
 RA Raport C.J., Gosling J., Schweichart V.L., Gray P.W., Charo I.F.;
 RT "Molecular cloning and functional characterization of a novel human
 RT CC chemokine receptor (CCR5) for RANTES, MIP-1beta, and MIP-1alpha.";
 RL J. Biol. Chem. 271:17161-17166(1996).
 RN [3]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=96295970; PubMed=8699119;
 RA Combadiere C., Ahuja S.K., Tiffany H.L., Murphy P.M.;
 RT "Cloning and functional expression of CC CKR5, a human monocyte CC
 RT chemokine receptor selective for MIP-1(alpha), MIP-1(beta), and
 RT RANTES.";
 RL J. Leukoc. Biol. 60:147-152(1996).
 RN [4]
 RP SEQUENCE FROM N.A.
 RA McCombie W.R., Wilson R., Chen E., Gibbs R., Zuo L., Johnson D.,
 RA Nhan M., Parnell L., Dedhia N., Ansari A., Mardis E., Schutz K.,
 RA Gnoj L., la Bastide M., Kaplan N., Greco T., Touchman J.,
 RA Muzny D., Chen C.N., Evans C., Fitzgerald M., See L.H., Tang M.,
 RA Porcel B.M., Dragan Y., Giacalone J., Pae A., Powell E.,
 RA Solinsky K.A., Desilva U., Diaz-Perez S., Zhou X., Yu Y.,
 RA Watanabe M., Doggett N., Garcia D., Sagripanti J.L.;
 RL Submitted (MAY-1997) to the EMBL/GenBank/DDBJ databases.
 RN [5]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=98001387; PubMed=9343222;
 RA Kuhmann S.E., Platt E.J., Kozak S.L., Kabat D.;
 RT "Polymorphisms in the CCR5 genes of African green monkeys and mice
 RT implicate specific amino acids in infections by simian and human
 RT immunodeficiency viruses.";
 RL J. Virol. 71:8642-8656(1997).
 RN [6]
 RP SEQUENCE FROM N.A., AND POLYMORPHISMS.
 RX MEDLINE=98022612; PubMed=9359654;
 RA Zhang L., Carruthers C.D., He T., Huang Y., Cao Y., Wang G., Hahn B.,
 RA Ho D.D.;
 RT "HIV type 1 subtypes, coreceptor usage, and CCR5 polymorphism.";
 RL AIDS Res. Hum. Retroviruses 13:1357-1366(1997).
 RN [7]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=98049523; PubMed=9388201;
 RA Mummidi S., Ahuja S.S., McDaniel B.L., Ahuja S.K.;
 RT "The human CC chemokine receptor 5 (CCR5) gene. Multiple transcripts
 RT with 5'-end heterogeneity, dual promoter usage, and evidence for
 RT polymorphisms within the regulatory regions and noncoding exons.";
 RL J. Biol. Chem. 272:30662-30671(1997).
 RN [8]
 RP SEQUENCE FROM N.A., AND VARIANT ARG-178.
 RA Magierowska M., Barre-Sinoussi F., Issafras H., Theodorou I.,
 RA Debre P.;
 RL Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.
 RN [9]
 RP SEQUENCE FROM N.A.
 RA Kopatz S.A., Aronstam R.S., Sharma S.V.;
 RT "cDNA clones of human proteins involved in signal transduction
 RT sequenced by the Guthrie cDNA resource center (www.cdna.org).";

RL Submitted (JAN-2003) to the EMBL/GenBank/DDBJ databases.
 RN [10]
 RP CHARACTERIZATION OF ITS HIV-1 RECEPTOR FUNCTION.
 RX MEDLINE=96260017; PubMed=8649511;
 RA Deng H., Liu R., Ellmeier W., Choe S., Unutmaz D., Burkhart M.,
 RA di Marzio P., Marmon S., Sutton R.E., Hill C.M., Davis C.B.,
 RA Peiper S.C., Schall T.J., Littman D.R., Landau N.R.;
 RT "Identification of a major co-receptor for primary isolates of
 RT HIV-1.";
 RL Nature 381:661-666(1996).
 RN [11]
 RP CHARACTERIZATION OF ITS HIV-1 RECEPTOR FUNCTION.
 RX MEDLINE=96260018; PubMed=8649512;
 RA Dragic T., Litwin V., Allaway G.P., Martin S.R., Huang Y.,
 RA Nagashima K.A., Cayanan C., Maddon P.J., Koup R.A., Moore J.P.,
 RA Paxton W.A.;
 RT "HIV-1 entry into CD4+ cells is mediated by the chemokine receptor
 RT CC-CKR-5.";
 RL Nature 381:667-673(1996).
 RN [12]
 RP SULFATION.
 RX MEDLINE=99189752; PubMed=10089882;
 RA Farzan M., Mirzabekov T., Kolchinsky P., Wyatt R., Cayabyab M.,
 RA Gerard N.P., Gerard C., Sodroski J., Choe H.;
 RT "Tyrosine sulfation of the amino terminus of CCR5 facilitates HIV-1
 RT entry.";
 RL Cell 96:667-676(1999).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and rantes and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation. Acts as co-receptor with CD4 for primary non-
 CC syncytium-inducing strains (NSI) (macrophage-tropic) of HIV-1
 CC virus. It promotes Env-mediated fusion of the virus.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- TISSUE SPECIFICITY: Found in promyelocytic cells.
 CC -!- PTM: Sulfation contributes to the efficiency of HIV-1 entry.
 CC -!- PTM: Modified by O-linked glycosylation, but not by N-linked
 CC glycosylation.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; X91492; CAA62796.1; -.
 DR EMBL; U54994; AAC50598.1; -.
 DR EMBL; U57840; AAB17071.1; -.
 DR EMBL; U95626; AAB57793.1; -.
 DR EMBL; U83326; AAC51797.1; -.
 DR EMBL; AF011500; AAB65700.1; -.
 DR EMBL; AF011501; AAB65701.1; -.
 DR EMBL; AF011502; AAB65702.1; -.

DR EMBL; AF011503; AAB65703.1; -.
 DR EMBL; AF011505; AAB65705.1; -.
 DR EMBL; AF011506; AAB65706.1; -.
 DR EMBL; AF011507; AAB65707.1; -.
 DR EMBL; AF011508; AAB65708.1; -.
 DR EMBL; AF011509; AAB65709.1; -.
 DR EMBL; AF011510; AAB65710.1; -.
 DR EMBL; AF011511; AAB65711.1; -.
 DR EMBL; AF011512; AAB65712.1; -.
 DR EMBL; AF011513; AAB65713.1; -.
 DR EMBL; AF011514; AAB65714.1; -.
 DR EMBL; AF011515; AAB65715.1; -.
 DR EMBL; AF011516; AAB65716.1; -.
 DR EMBL; AF011517; AAB65717.1; -.
 DR EMBL; AF011518; AAB65718.1; -.
 DR EMBL; AF011519; AAB65719.1; -.
 DR EMBL; AF011520; AAB65720.1; -.
 DR EMBL; AF011521; AAB65721.1; -.
 DR EMBL; AF011522; AAB65722.1; -.
 DR EMBL; AF011523; AAB65723.1; -.
 DR EMBL; AF011524; AAB65724.1; -.
 DR EMBL; AF011525; AAB65725.1; -.
 DR EMBL; AF011526; AAB65726.1; -.
 DR EMBL; AF011527; AAB65727.1; -.
 DR EMBL; AF011528; AAB65728.1; -.
 DR EMBL; AF011529; AAB65729.1; -.
 DR EMBL; AF011530; AAB65730.1; -.
 DR EMBL; AF011531; AAB65731.1; -.
 DR EMBL; AF011532; AAB65732.1; -.
 DR EMBL; AF011533; AAB65733.1; -.
 DR EMBL; AF011534; AAB65734.1; -.
 DR EMBL; AF011535; AAB65735.1; -.
 DR EMBL; AF011536; AAB65736.1; -.
 DR EMBL; AF011537; AAB65737.1; -.
 DR EMBL; AF031237; AAB94735.1; -.
 DR EMBL; AF052539; AAD18131.1; -.
 DR EMBL; AY221093; AAO65971.1; -.
 DR Genew; HGNC:1606; CCR5.
 DR MIM; 601373; -.
 DR GO; GO:0005768; C:endosome; TAS.
 DR GO; GO:0005887; C:integral to plasma membrane; TAS.
 DR GO; GO:0016493; F:C-C chemokine receptor activity; NAS.
 DR GO; GO:0015026; F:coreceptor activity; TAS.
 DR GO; GO:0007267; P:cell-cell signaling; TAS.
 DR GO; GO:0006968; P:cellular defense response; TAS.
 DR GO; GO:0006935; P:chemotaxis; TAS.
 DR GO; GO:0007204; P:cytosolic calcium ion concentration elevation; TAS.
 DR GO; GO:0007186; P:G-protein coupled receptor protein signalin. . .; TAS.
 DR GO; GO:0006954; P:inflammatory response; TAS.
 DR GO; GO:0007125; P:invasive growth; TAS.
 DR GO; GO:0007203; P:phosphatidylinositol-4,5-bisphosphate hydro. . .; TAS.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS0262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation;

KW	Polymorphism.			
FT	DOMAIN	1	30	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	31	58	1 (POTENTIAL).
FT	DOMAIN	59	68	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	69	89	2 (POTENTIAL).
FT	DOMAIN	90	102	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	103	124	3 (POTENTIAL).

Query Match 42.9%; Score 379; DB 1; Length 352;
 Best Local Similarity 35.0%; Pred. No. 2.2e-29;
 Matches 99; Conservative 7; Mismatches 27; Indels 150; Gaps 7;

Qy	20	MDYQVSSPIYDINYYTSEPCQKINVKQIAA-----	49
Db	1	MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR	60
Qy	50	-----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG---	83
		: :	
Db	61	LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF	118
Qy	84	-----LCTRSQKEGLHYTC	97
		:	
Db	119	IILLTIDRYLAVVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC	178
Qy	98	SSHFPYSQYQFWKNFQTLKI-----HQRVHGG-----	124
		:	
Db	179	SSHFPYSQYQFWKNFQTLKIVILGLVLPLLVMVICYSGILKTLLRCRNEKKRHRAVRLIF	238
Qy	125	-----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV	154
		:	
Db	239	TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV	281

RESULT 2

CKR5_PANTR

ID CKR5_PANTR STANDARD; PRT; 352 AA.
 AC P56440; O02778;
 DT 15-JUL-1998 (Rel. 36, Created)
 DT 15-JUL-1998 (Rel. 36, Last sequence update)
 DT 16-OCT-2001 (Rel. 40, Last annotation update)
 DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).
 GN CCR5 OR CMKBR5.
 OS Pan troglodytes (Chimpanzee).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Pan.
 OX NCBI_TaxID=9598;
 RN [1]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=97268687; PubMed=9108095;
 RA Edinger A.L., Amedee A., Miller K., Doranz B.J., Endres M.,
 RA Sharron M., Samson M., Lu Z.H., Clements J.E., Murphey-Corb M.,
 RA Peiper S.C., Parmentier M., Broder C.C., Doms R.W.;
 RT "Differential utilization of CCR5 by macrophage and T cell tropic
 RT simian immunodeficiency virus strains."
 RL Proc. Natl. Acad. Sci. U.S.A. 94:4005-4010(1997).
 RN [2]
 RP SEQUENCE FROM N.A.

RA Zimmerman P.A., Buckler-White A., Alkhatib G.;
 RL Submitted (MAR-1997) to the EMBL/GenBank/DDBJ databases.
 RN [3]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=98022612; PubMed=9359654;
 RA Zhang L., Carruthers C.D., He T., Huang Y., Cao Y., Wang G., Hahn B.,
 RA Ho D.D.;
 RT "HIV type 1 subtypes, coreceptor usage, and CCR5 polymorphism.";
 RL AIDS Res. Hum. Retroviruses 13:1357-1366(1997).
 RN [4]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=97426118; PubMed=9282822;
 RA Zacharova V., Zachar V., Goustin A.S.;
 RT "Sequence of chemokine receptor gene CCR5 in chimpanzees, a natural
 RT HIV type 1 host.";
 RL AIDS Res. Hum. Retroviruses 13:1159-1161(1997).
 RN [5]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=98090115; PubMed=9430250;
 RA Pretet J.-L., Zerbib A., Girard M., Guillet J.-G., Butor C.;
 RT "Chimpanzee CXCR4 and CCR5 act as coreceptors for HIV type 1.";
 RL AIDS Res. Hum. Retroviruses 13:1583-1587(1997).
 RN [6]
 RP SEQUENCE FROM N.A.
 RA Zhang Y., Ryder O.A., Zhang Y.;
 RT "Sequence comparison of the CCR5 gene in primates and primate
 RT phylogeny.";
 RL Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF005663; AAB62557.1; -.
 DR EMBL; U94329; AAB58446.1; -.
 DR EMBL; AF011542; AAB65742.1; -.
 DR EMBL; U97666; AAC51670.1; -.
 DR EMBL; AF011540; AAB65740.1; -.
 DR EMBL; U89797; AAC03717.1; -.
 DR EMBL; AF177894; AAK43377.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.

FT	DOMAIN	1	30	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	31	58	1 (POTENTIAL).
FT	DOMAIN	59	68	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	69	89	2 (POTENTIAL).
FT	DOMAIN	90	102	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	103	124	3 (POTENTIAL).
FT	DOMAIN	125	141	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	142	166	4 (POTENTIAL).
FT	DOMAIN	167	198	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	199	218	5 (POTENTIAL).
FT	DOMAIN	219	235	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	236	260	6 (POTENTIAL).
FT	DOMAIN	261	277	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	278	301	7 (POTENTIAL).
FT	DOMAIN	302	352	CYTOPLASMIC (POTENTIAL).
FT	DISULFID	101	178	BY SIMILARITY.
FT	MOD_RES	3	3	SULFATION (BY SIMILARITY).
FT	MOD_RES	10	10	SULFATION (BY SIMILARITY).
FT	MOD_RES	14	14	SULFATION (BY SIMILARITY).
FT	MOD_RES	15	15	SULFATION (BY SIMILARITY).
FT	CARBOHYD	268	268	N-LINKED (GLCNAC. . .) (POTENTIAL).
FT	CONFLICT	123	123	T -> S (IN REF. 1).
SO	SEQUENCE	352 AA;	40539 MW;	4A33E698B80FE34C CRC64;

```

Qy      20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      |||||||||||:|||||||||||||||
Db      1 MDYQVSSPIYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | | | | | | | | | | | | | : | : : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : |||||||||
Db      119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      ||||||||||| | | | | | | | | | : : |
Db      179 SSHFPYSQYQFWKNFQTLKIVILGLVLPPLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238

Qy      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
      | | | | | | | | | | | | | | : |
Db      239 TIMIVYFLFWAPYNIVLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 3

ID CKR5 HYLSY STANDARD; PRT; 352 AA.

DT 15-MAR-2004 (Rel. 43, Created)

DT 15-MAR-2004 (Rel. 43, Last annotation update)

GN CCR5 OR CMKBR5.
 OS Hylobates syndactylus (Siamang) (Symphalangus syndactylus).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Hylobatidae; Hylobates.
 OX NCBI_TaxID=9590;
 RN [1]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=99416438; PubMed=10486970;
 RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;
 RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";
 RL Mol. Biol. Evol. 16:1145-1154(1999).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF177884; AAK43367.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40508 MW; F4F64B3AD5AF658A CRC64;

Query Match 41.7%; Score 369; DB 1; Length 352;
 Best Local Similarity 34.3%; Pred. No. 2e-28;

Matches 97; Conservative 8; Mismatches 28; Indels 150; Gaps 7;

```

Qy      20 MDYQVSSPIYDINYTSEPCQKINVKQIAA----- 49
      ||||| |||:|||||
Db      1 MDYQVSSPTYDIDYITSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60

Qy      50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
      | | ||||| | : | : |
Db      61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy      84 -----LCTRSQKEGLHYTC 97
      : |||||
Db     119 IILLTIDRYLAIVHAVFALKARTVTFGVTVSVITWVAVFASLPGIIFTRSQKEGLHYTC 178

Qy      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      ||||| ||||| : : |
Db     179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLRLCRNEKKRHRVRLIF 238

Qy     125 -----GGSYKCGLC----QEFGGLNNCSSSNRLDGHQRV 154
      | | ||||| : |
Db     239 TIMIVYFLFWAPYNIVLLNTFQEFGGLNNCSSSNRLDQAMQV 281

```

RESULT 4

CKR5_PONPY

ID CKR5_PONPY STANDARD; PRT; 352 AA.

AC O97881;

DT 30-MAY-2000 (Rel. 39, Created)

DT 30-MAY-2000 (Rel. 39, Last sequence update)

DT 28-FEB-2003 (Rel. 41, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).

GN CCR5 OR CMKBR5.

OS Pongo pygmaeus (Orangutan).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Pongo.

OX NCBI_TaxID=9600;

RN [1]

RP SEQUENCE FROM N.A.

RX MEDLINE=99416438; PubMed=10486970;

RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;

RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";

RL Mol. Biol. Evol. 16:1145-1154(1999).

CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.

CC -!- SUBCELLULAR LOCATION: Integral membrane protein.

CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.

CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).

CC -----
 DR EMBL; AF075446; AAD19858.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40527 MW; F4E2F47135AF658A CRC64;

Query Match 41.7%; Score 369; DB 1; Length 352;
 Best Local Similarity 34.3%; Pred. No. 2e-28;
 Matches 97; Conservative 8; Mismatches 28; Indels 150; Gaps 7;

QY 20 MDYQVSSPIYDINYYTSEPCQKINVQIAA----- 49
 ||||| :|||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
 QY 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 QY 84 -----LCTRSQKEGLHYTC 97
 : |||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178
 QY 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| :|
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRAVRLIF 238
 QY 125 -----GGSYKCGLC-----QEFGGLNNCSSSNRLDGHQRV 154
 | | ||||| :|
 Db 239 TIMIVYFLFWAPYNIIVLLNTFQEFGGLNNCSSSNRLDQAMQV 281

RESULT 5
 CKR5_PAPHA

ID CKR5_PAPHA STANDARD; PRT; 352 AA.
 AC P56441;
 DT 15-JUL-1998 (Rel. 36, Created)
 DT 15-JUL-1998 (Rel. 36, Last sequence update)
 DT 16-OCT-2001 (Rel. 40, Last annotation update)
 DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).
 GN CCR5 OR CMKBR5.
 OS Papio hamadryas (Hamadryas baboon), and
 OS Papio anubis (Olive baboon).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; Papio.
 OX NCBI_TaxID=9557, 9555;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC SPECIES=P.hamadryas;
 RX MEDLINE=97268687; PubMed=9108095;
 RA Edinger A.L., Amedee A., Miller K., Doranz B.J., Endres M.,
 RA Sharron M., Samson M., Lu Z.H., Clements J.E., Murphey-Corb M.,
 RA Peiper S.C., Parmentier M., Broder C.C., Doms R.W.;
 RT "Differential utilization of CCR5 by macrophage and T cell tropic
 RT simian immunodeficiency virus strains.";
 RL Proc. Natl. Acad. Sci. U.S.A. 94:4005-4010(1997).
 RN [2]
 RP SEQUENCE FROM N.A.
 RC SPECIES=P.hamadryas;
 RX MEDLINE=99210133; PubMed=10195758;
 RA Saksena N.K., Wang B., Novembre F.J., Bolton W., Smit T.K., Lal R.B.;
 RT "Species-specific changes in the CCR5 gene from African and Asian
 RT nonhuman primates.";
 RL AIDS Res. Hum. Retroviruses 15:479-483(1999).
 RN [3]
 RP SEQUENCE FROM N.A.
 RC SPECIES=P.anubis;
 RA Benton P.A., Timanus D.K., Shearer M.H., Lee D.R., Kennedy R.C.;
 RL Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF005658; AAB62552.1; -.
 DR EMBL; AF105287; AAD20556.1; -.
 DR EMBL; AF105288; AAD20557.1; -.
 DR EMBL; AF105289; AAD20558.1; -.
 DR EMBL; AF105290; AAD20559.1; -.

DR EMBL; AF023452; AAC63830.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 FT CARBOHYD 268 268 N-LINKED (GLCNAC. . .) (POTENTIAL).
 SQ SEQUENCE 352 AA; 40489 MW; 5E1504A9BA1FE8B2 CRC64;

Query Match 41.4%; Score 366; DB 1; Length 352;
 Best Local Similarity 33.9%; Pred. No. 3.9e-28;
 Matches 96; Conservative 9; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 Db 61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLRLCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFGGLNNCSSSNRLDGHQRV 154
 Db 239 TIMIVYFLFWAPYNIVLLNLTFFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 6
 CKR5_PYGBI

ID CKR5_PYGBI STANDARD; PRT; 352 AA.
 AC O97880;
 DT 30-MAY-2000 (Rel. 39, Created)
 DT 30-MAY-2000 (Rel. 39, Last sequence update)
 DT 28-FEB-2003 (Rel. 41, Last annotation update)
 DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).
 GN CCR5 OR CMKBR5.
 OS *Pygathrix bieti* (Black snub-nosed monkey) (*Rhinopithecus bieti*).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;
 OC *Pygathrix*.
 OX NCBI_TaxID=61621;
 RN [1]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=99416438; PubMed=10486970;
 RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;
 RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";
 RL Mol. Biol. Evol. 16:1145-1154(1999).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF075445; AAD19857.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.

FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40585 MW; 4366F142730F938F CRC64;

Query Match 41.3%; Score 365; DB 1; Length 352;
 Best Local Similarity 33.6%; Pred. No. 4.9e-28;
 Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||:|||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKVNKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG--- 83
 | | ||||| | : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQREGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| ||||| : : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGLKTLRLCRNEKKRHRVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFGFLNNCSSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLEWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQV 281

RESULT 7

CKR5_PYGNE

ID CKR5_PYGNE STANDARD; PRT; 352 AA.

AC O97882;

DT 30-MAY-2000 (Rel. 39, Created)

DT 30-MAY-2000 (Rel. 39, Last sequence update)

DT 28-FEB-2003 (Rel. 41, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).

GN CCR5 OR CMKBR5.

OS Pygathrix nemaeus (Dove langur).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;

OC Pygathrix.

OX NCBI_TaxID=54133;

RN [1]

RP SEQUENCE FROM N.A.

RX MEDLINE=99416438; PubMed=10486970;

RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;

RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";

RL Mol. Biol. Evol. 16:1145-1154(1999).

CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.

CC -!- SUBCELLULAR LOCATION: Integral membrane protein.

Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLIMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

Qy 125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154

Db 239 TIMIVYFLFWAPYNIV LLLNTFQEFFGLNNC SSSNRLDQAMQV 281

RESULT 8

CKR5_TRAFR

ID CKR5_TRAFR STANDARD; PRT; 352 AA.

AC O97878;

DT 30-MAY-2000 (Rel. 39, Created)

DT 30-MAY-2000 (Rel. 39, Last sequence update)

DT 28-FEB-2003 (Rel. 41, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).

GN CCR5 OR CMKBR5.

OS Trachypithecus francoisi (Francois' langur) (Indochinese langur).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;

OC Trachypithecus.

OX NCBI_TaxID=54180;

RN [1]

RP SEQUENCE FROM N.A.

RX MEDLINE=99416438; PubMed=10486970;

RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;

RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";

RL Mol. Biol. Evol. 16:1145-1154(1999).

CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,

CC MIP-1-beta and RANTES and subsequently transduces a signal by

CC increasing the intracellular calcium ions level. May play a role

CC in the control of granulocytic lineage proliferation or

CC differentiation.

CC -!- SUBCELLULAR LOCATION: Integral membrane protein.

CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.

CC -----

CC This SWISS-PROT entry is copyright. It is produced through a collaboration

CC between the Swiss Institute of Bioinformatics and the EMBL outstation -

CC the European Bioinformatics Institute. There are no restrictions on its

CC use by non-profit institutions as long as its content is in no way

CC modified and this statement is not removed. Usage by and for commercial

CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>

CC or send an email to license@isb-sib.ch).

CC -----

DR EMBL; AF075442; AAD19854.1; -.

DR InterPro; IPR000276; GPCR_Rhodpsn.

DR Pfam; PF00001; 7tm_1; 1.

DR PRINTS; PR00237; GPCRRHODOPSN.

DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.

DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.

KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.

FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).

FT TRANSMEM 31 58 1 (POTENTIAL).

FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).

FT TRANSMEM 69 89 2 (POTENTIAL).

FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).

FT TRANSMEM 103 124 3 (POTENTIAL).

FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).

FT	TRANSMEM	142	166	4 (POTENTIAL).
FT	DOMAIN	167	198	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	199	218	5 (POTENTIAL).
FT	DOMAIN	219	235	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	236	260	6 (POTENTIAL).
FT	DOMAIN	261	277	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	278	301	7 (POTENTIAL).
FT	DOMAIN	302	352	CYTOPLASMIC (POTENTIAL).
FT	DISULFID	101	178	BY SIMILARITY.
FT	MOD_RES	3	3	SULFATION (BY SIMILARITY).
FT	MOD_RES	10	10	SULFATION (BY SIMILARITY).
FT	MOD_RES	14	14	SULFATION (BY SIMILARITY).
FT	MOD_RES	15	15	SULFATION (BY SIMILARITY).
SQ	SEQUENCE	352 AA;	40496 MW;	4366F148C255938F CRC64;

Query Match 41.3%; Score 365; DB 1; Length 352;
 Best Local Similarity 33.6%; Pred. No. 4.9e-28;
 Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

Qy	20	MDYQVSSPIYDINYYTSEPCQKINVKQIAA-----	49
		: : : : : : : : :	
Db	1	MDYQVSSPTYDIDYYTSEPCQKVNKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR	60
Qy	50	-----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG---	83
		: : :	
Db	61	LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF	118
Qy	84	-----LCTRSQKEGLHYTC	97
		: :	
Db	119	IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQREGLHYTC	178
Qy	98	SSHFPYSQYQFWKNFQTLKI-----HQRVHGG-----	124
		:	
Db	179	SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYS GILKTL LRCRNEKKRHRAVRLIF	238
Qy	125	-----GGSYKCGLC-----QEFFGLNNC SSSNRLDGHQRV	154
		:	
Db	239	TIMIVYFLEWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV	281

RESULT 9

CKR5_TRAPH

ID CKR5_TRAPH STANDARD; PRT; 352 AA.

AC O97879;

DT 30-MAY-2000 (Rel. 39, Created)

DT 30-MAY-2000 (Rel. 39, Last sequence update)

DT 28-FEB-2003 (Rel. 41, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).

GN CCR5 OR CMKBR5.

OS Trachypithecus phayrei (Phayre's leaf monkey).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae; Colobinae;

OC Trachypithecus.

OX NCBI_TaxID=61618;

RN [1]

RP SEQUENCE FROM N.A.

RX MEDLINE=99416438; PubMed=10486970;

RA Zhang Y.-W., Ryder O.A., Zhang Y.-P.;
 RT "Sequence evolution of the CCR5 chemokine receptor gene in primates.";
 RL Mol. Biol. Evol. 16:1145-1154(1999).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF075443; AAD19855.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40509 MW; 4366F148D3A5938F CRC64;

Query Match 41.3%; Score 365; DB 1; Length 352;
 Best Local Similarity 33.6%; Pred. No. 4.9e-28;
 Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

QY 20 MDYQVSSPIYDINYTSEPCQKINVKQIAA----- 49
 ||||| |||:|||||||:|||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKVNKQIAARLLPLYSLVFIFGFVGNILVVLILINCKR 60
 QY 50 -----YKGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83

```

Db          61 LKSMTDIYLLNLAIISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
Qy          84 -----LCTRSQKEGLHYTC 97
              : ||||:|||||
Db          119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIIFTRSQREGLHYTC 178
Qy          98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
              |||||:|
Db          179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTLLRCRNEKKRHRVRLIF 238
Qy          125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
              | | |||||:|
Db          239 TIMIVYFLWAPYNIVLLNTFOEFFGLNNCSSSNRLDQAMQV 281

```

DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40515 MW; D0E6FCB9FE5EAC84 CRC64;

Query Match 41.2%; Score 364; DB 1; Length 352;
 Best Local Similarity 33.9%; Pred. No. 6.1e-28;
 Matches 96; Conservative 8; Mismatches 29; Indels 150; Gaps 7;

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| : |||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKTNVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR 60
 Qy 50 -----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| : | : |
 Db 61 LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 Qy 84 -----LCTRSQKEGLHYTC 97
 : |||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQKEGLHYTC 178
 Qy 98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
 ||||| : |
 Db 179 SSHFPYSQYQFWKNFQTLKIVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238
 Qy 125 -----GGSYKCGLC----QEFFGLNNC SSSNRLDGHQRV 154
 | | ||||| : |
 Db 239 TIMIVYFLFWAPYNIVLLLNTFQEFFGLNNC SSSNRLDQAMQV 281

RESULT 11

CKR5_MACMU

ID CKR5_MACMU STANDARD; PRT; 352 AA.
 AC P79436; 002746;
 DT 01-NOV-1997 (Rel. 35, Created)
 DT 01-NOV-1997 (Rel. 35, Last sequence update)

DT 28-FEB-2003 (Rel. 41, Last annotation update)
 DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).
 GN CCR5 OR CMKBR5.
 OS *Macaca mulatta* (Rhesus macaque),
 OS *Macaca fascicularis* (Crab eating macaque) (*Cynomolgus* monkey), and
 OS *Macaca nemestrina* (Pig-tailed macaque).
 OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;
 OC Cercopithecinae; *Macaca*.
 OX NCBI_TaxID=9544, 9541, 9545;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC SPECIES=*M.mulatta*;
 RX MEDLINE=97184592; PubMed=9032394;
 RA Marcon L., Choe H., Martin K.A., Farzan M., Ponath P.D., Wu L.,
 RA Newman W., Gerard N., Gerard C., Sodroski J.;
 RT "Utilization of C-C chemokine receptor 5 by the envelope
 RT glycoproteins of a pathogenic simian immunodeficiency virus,
 RT SIVmac239.";
 RL J. Virol. 71:2522-2527(1997).
 RN [2]
 RP SEQUENCE FROM N.A.
 RC SPECIES=*M.mulatta*; STRAIN=Indian macaque;
 RX MEDLINE=97213934; PubMed=9060623;
 RA Chen Z., Zhou P., Ho D.D., Landau N.R., Marx P.A.;
 RT "Genetically divergent strains of simian immunodeficiency virus use
 RT CCR5 as a coreceptor for entry.";
 RL J. Virol. 71:2705-2714(1997).
 RN [3]
 RP SEQUENCE FROM N.A.
 RC SPECIES=*M.mulatta*;
 RX MEDLINE=21354176; PubMed=11461684;
 RA Margulies B.J., Hauer D.A., Clements J.E.;
 RT "Identification and comparison of eleven rhesus macaque chemokine
 RT receptors.";
 RL AIDS Res. Hum. Retroviruses 17:981-986(2001).
 RN [4]
 RP SEQUENCE FROM N.A.
 RC SPECIES=*M.mulatta*, *M.fascicularis*, and *M.nemestrina*;
 RX MEDLINE=97268687; PubMed=9108095;
 RA Edinger A.L., Amedee A., Miller K., Doranz B.J., Endres M.,
 RA Sharron M., Samson M., Lu Z.H., Clements J.E., Murphey-Corb M.,
 RA Peiper S.C., Parmentier M., Broder C.C., Doms R.W.;
 RT "Differential utilization of CCR5 by macrophage and T cell tropic
 RT simian immunodeficiency virus strains.";
 RL Proc. Natl. Acad. Sci. U.S.A. 94:4005-4010(1997).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its

CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).

CC -----
 DR EMBL; U77672; AAC51109.1; -.
 DR EMBL; U73739; AAC51158.1; -.
 DR EMBL; U96762; AAC34132.1; -.
 DR EMBL; AF005660; AAB62554.1; -.
 DR EMBL; AF005661; AAB62555.1; -.
 DR EMBL; AF005662; AAB62556.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 FT CARBOHYD 268 268 N-LINKED (GLCNAC. . .) (POTENTIAL).
 FT CONFLICT 241 241 M -> I (IN REF. 3).
 FT CONFLICT 292 292 I -> M (IN REF. 3).
 SQ SEQUENCE 352 AA; 40507 MW; 58B96C85909FACB2 CRC64;

Query Match 41.1%; Score 363; DB 1; Length 352;
 Best Local Similarity 33.6%; Pred. No. 7.6e-28;
 Matches 95; Conservative 10; Mismatches 28; Indels 150; Gaps 7;

QY 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| :|||||||
 Db 1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60
 QY 50 -----YKCGLC-----AAQWDFGNTMCQHQRVHGHHSYKCG--- 83
 | | ||||| : | : |
 Db 61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118
 QY 84 -----LCTRSQKEGLHYTC 97
 : |||:|||||
 Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGIIFTRSQREGHYTC 178

```

QY      98 SSHFPYSQYQFWKNFQTLKI-----HQRVHGG----- 124
      |||||||||||||||||:
Db      179 SSHFPYSQYQFWKNFQTLKMVILGLVLP LLVMVICYSGILKTL LRCRNEKKRHRAVRLIF 238

QY      125 -----GGSYKCGLC----QEFFGLNNCSSSNRLDGHQRV 154
      | | |||||||||||||:|
Db      239 TIMIVYFLFWAPYNIVLLNTFQEFFGLNNCSSSNRLDQAMQV 281

```

RESULT 12

CKR5_HYLML

```

ID      CKR5_HYLML      STANDARD;      PRT;      352 AA.
AC      Q95NC0;
DT      15-MAR-2004 (Rel. 43, Created)
DT      15-MAR-2004 (Rel. 43, Last sequence update)
DT      15-MAR-2004 (Rel. 43, Last annotation update)
DE      C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).
GN      CCR5 OR CMKBR5.
OS      Hylobates moloch (Silvery gibbon).
OC      Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC      Mammalia; Eutheria; Primates; Catarrhini; Hylobatidae; Hylobates.
OX      NCBI_TaxID=81572;
RN      [1]
RP      SEQUENCE FROM N.A.
RX      MEDLINE=99416438; PubMed=10486970;
RA      Zhang Y.-W., Ryder O.A., Zhang Y.-P.;
RT      "Sequence evolution of the CCR5 chemokine receptor gene in primates.";
RL      Mol. Biol. Evol. 16:1145-1154(1999).
CC      -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
CC      MIP-1-beta and RANTES and subsequently transduces a signal by
CC      increasing the intracellular calcium ions level. May play a role
CC      in the control of granulocytic lineage proliferation or
CC      differentiation.
CC      -!- SUBCELLULAR LOCATION: Integral membrane protein.
CC      -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
CC      -----
CC      This SWISS-PROT entry is copyright. It is produced through a collaboration
CC      between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC      the European Bioinformatics Institute. There are no restrictions on its
CC      use by non-profit institutions as long as its content is in no way
CC      modified and this statement is not removed. Usage by and for commercial
CC      entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC      or send an email to license@isb-sib.ch).
CC      -----
DR      EMBL; AF177899; AAK43382.1; -.
DR      InterPro; IPR000276; GPCR_Rhodpsn.
DR      Pfam; PF00001; 7tm_1; 1.
DR      PRINTS; PR00237; GPCRRHODOPSN.
DR      PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR      PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
KW      G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
FT      DOMAIN      1      30      EXTRACELLULAR (POTENTIAL).
FT      TRANSMEM      31      58      1 (POTENTIAL).
FT      DOMAIN      59      68      CYTOPLASMIC (POTENTIAL).
FT      TRANSMEM      69      89      2 (POTENTIAL).
FT      DOMAIN      90      102      EXTRACELLULAR (POTENTIAL).

```

FT	TRANSMEM	103	124	3 (POTENTIAL).
FT	DOMAIN	125	141	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	142	166	4 (POTENTIAL).
FT	DOMAIN	167	198	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	199	218	5 (POTENTIAL).
FT	DOMAIN	219	235	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	236	260	6 (POTENTIAL).
FT	DOMAIN	261	277	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	278	301	7 (POTENTIAL).
FT	DOMAIN	302	352	CYTOPLASMIC (POTENTIAL).
FT	DISULFID	101	178	BY SIMILARITY.
FT	MOD_RES	3	3	SULFATION (BY SIMILARITY).
FT	MOD_RES	10	10	SULFATION (BY SIMILARITY).
FT	MOD_RES	14	14	SULFATION (BY SIMILARITY).
SQ	SEQUENCE	352 AA;	40436 MW;	9623CA98340CF274 CRC64;

Query Match 41.0%; Score 362; DB 1; Length 352;

Best Local Similarity 33.9%; Pred. No. 9.5e-28;

Matches 96; Conservative 8; Mismatches 29; Indels 150; Gaps 7;

Qy	20	MDYQVSSPIYDINYYTSEPCQKINVKQIAA-----	49
		:	
Db	1	MDYQVSSPTYDIDYYTSGPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKR	60
Qy	50	-----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG---	83
		: :	
Db	61	LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF	118
Qy	84	-----LCTRSQKEGLHYTC	97
		:	
Db	119	IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVAVFASLPGLIIFTRSQKEGLHYTC	178
Qy	98	SSHFPYSQYQFWKNFQTLKI-----HQRVHGG-----	124
		: :	
Db	179	SSHFPYSQYQFWKNFQTLKIVILGLVLPPLVMVICYSGLKTLRLCRNEKKRHRRAVRLIF	238
Qy	125	-----GGSYKCGLC---QEFGGLNNCSSSNRLDGHQRV	154
		:	
Db	239	TIMIVYFLFWAPYNIVLLLNTFQEFGGLNNCSSSNRLDQAMQV	281

RESULT 13

CKR5_CERP

ID CKR5_CERP STANDARD; PRT; 352 AA.

AC Q9TV42;

DT 15-MAR-2004 (Rel. 43, Created)

DT 15-MAR-2004 (Rel. 43, Last sequence update)

DT 15-MAR-2004 (Rel. 43, Last annotation update)

DE C-C chemokine receptor type 5 (C-C CKR-5) (CC-CKR-5) (CCR-5) (CCR5).

GN CCR5 OR CMKBR5.

OS Cercopithecus pygerythrus (Vervet monkey).

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

OC Mammalia; Eutheria; Primates; Catarrhini; Cercopithecidae;

OC Cercopithecinae; Cercopithecus.

OX NCBI_TaxID=60710;

RN [1]

RP SEQUENCE FROM N.A.

RC STRAIN=100;
 RX MEDLINE=99335215; PubMed=10408730;
 RA Muller-Trutwin M.C., Corbet S., Hansen J., Georges-Courbot M.-C.,
 RA Diop O., Rigoulet J., Barre-Sinoussi F., Fomsgaard A.;
 RT "Mutations in CCR5-coding sequences are not associated with SIV
 RT carrier status in African nonhuman primates.";
 RL AIDS Res. Hum. Retroviruses 15:931-939(1999).
 CC -!- FUNCTION: Receptor for a C-C type chemokine. Binds to MIP-1-alpha,
 CC MIP-1-beta and RANTES and subsequently transduces a signal by
 CC increasing the intracellular calcium ions level. May play a role
 CC in the control of granulocytic lineage proliferation or
 CC differentiation.
 CC -!- SUBCELLULAR LOCATION: Integral membrane protein.
 CC -!- SIMILARITY: Belongs to family 1 of G-protein coupled receptors.
 CC -----
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC -----
 DR EMBL; AF035222; AAD44015.1; -.
 DR InterPro; IPR000276; GPCR_Rhodpsn.
 DR Pfam; PF00001; 7tm_1; 1.
 DR PRINTS; PR00237; GPCRRHODOPSN.
 DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
 DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
 KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
 FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 31 58 1 (POTENTIAL).
 FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 69 89 2 (POTENTIAL).
 FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 103 124 3 (POTENTIAL).
 FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 142 166 4 (POTENTIAL).
 FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 199 218 5 (POTENTIAL).
 FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
 FT TRANSMEM 236 260 6 (POTENTIAL).
 FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
 FT TRANSMEM 278 301 7 (POTENTIAL).
 FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
 FT DISULFID 101 178 BY SIMILARITY.
 FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
 FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
 FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
 SQ SEQUENCE 352 AA; 40558 MW; EF17D67CBCCC3DB0 CRC64;

Query Match 40.8%; Score 361; DB 1; Length 352;
 Best Local Similarity 33.6%; Pred. No. 1.2e-27;
 Matches 95; Conservative 9; Mismatches 29; Indels 150; Gaps 7;

QY 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
 ||||| :|||||


```

CC -----
DR EMBL; AF051902; AAC39830.1; -.
DR EMBL; AF051903; AAC39831.1; -.
DR EMBL; AF051904; AAC39832.1; -.
DR EMBL; AF051905; AAC39833.1; -.
DR InterPro; IPR000276; GPCR_Rhodpsn.
DR Pfam; PF00001; 7tm_1; 1.
DR PRINTS; PR00237; GPCRRHODOPSN.
DR PROSITE; PS00237; G_PROTEIN_RECEP_F1_1; 1.
DR PROSITE; PS50262; G_PROTEIN_RECEP_F1_2; 1.
KW G-protein coupled receptor; Transmembrane; Glycoprotein; Sulfation.
FT DOMAIN 1 30 EXTRACELLULAR (POTENTIAL).
FT TRANSMEM 31 58 1 (POTENTIAL).
FT DOMAIN 59 68 CYTOPLASMIC (POTENTIAL).
FT TRANSMEM 69 89 2 (POTENTIAL).
FT DOMAIN 90 102 EXTRACELLULAR (POTENTIAL).
FT TRANSMEM 103 124 3 (POTENTIAL).
FT DOMAIN 125 141 CYTOPLASMIC (POTENTIAL).
FT TRANSMEM 142 166 4 (POTENTIAL).
FT DOMAIN 167 198 EXTRACELLULAR (POTENTIAL).
FT TRANSMEM 199 218 5 (POTENTIAL).
FT DOMAIN 219 235 CYTOPLASMIC (POTENTIAL).
FT TRANSMEM 236 260 6 (POTENTIAL).
FT DOMAIN 261 277 EXTRACELLULAR (POTENTIAL).
FT TRANSMEM 278 301 7 (POTENTIAL).
FT DOMAIN 302 352 CYTOPLASMIC (POTENTIAL).
FT DISULFID 101 178 BY SIMILARITY.
FT MOD_RES 3 3 SULFATION (BY SIMILARITY).
FT MOD_RES 10 10 SULFATION (BY SIMILARITY).
FT MOD_RES 14 14 SULFATION (BY SIMILARITY).
FT MOD_RES 15 15 SULFATION (BY SIMILARITY).
FT VARIANT 2 2 D -> E (IN ISOLATE 087).
FT VARIANT 3 3 Y -> D (IN ISOLATE 079).
FT VARIANT 25 25 V -> G (IN ISOLATE 087).
FT VARIANT 100 100 M -> K (IN ISOLATE 079).
FT VARIANT 107 107 L -> V (IN ISOLATE 089).
FT VARIANT 134 134 V -> G (IN ISOLATE 079).
FT VARIANT 146 146 V -> L (IN ISOLATES 085 AND 089).
FT VARIANT 340 340 T -> I (IN ISOLATE 079).
SQ SEQUENCE 352 AA; 40489 MW; 20A196E2D47E49CA CRC64;

```

```

Query Match 40.8%; Score 361; DB 1; Length 352;
Best Local Similarity 33.6%; Pred. No. 1.2e-27;
Matches 95; Conservative 9; Mismatches 29; Indels 150; Gaps 7;

```

```

Qy 20 MDYQVSSPIYDINYYTSEPCQKINVKQIAA----- 49
      ||||| |||:|||||
Db 1 MDYQVSSPTYDIDYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNILVVLILINCKR 60

Qy 50 -----YKGLC-----AAAQWDFGNTMCQHQRVHGHHSYKCG--- 83
      | | ||||| | : | : |
Db 61 LKSMTDIYLLNLAISDLLFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF 118

Qy 84 -----LCTRSQKEGLHYTC 97
      : |||:|||||
Db 119 IILLTIDRYLAIVHAVFALKARTVTFGVVTSVITWVVAVFASLPGIIFTRSQREGLHYTC 178

```

FT TRANSMEM 103 124 3 (POTENTIAL).

FT	DOMAIN	125	141	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	142	166	4 (POTENTIAL).
FT	DOMAIN	167	198	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	199	218	5 (POTENTIAL).
FT	DOMAIN	219	235	CYTOPLASMIC (POTENTIAL).
FT	TRANSMEM	236	260	6 (POTENTIAL).
FT	DOMAIN	261	277	EXTRACELLULAR (POTENTIAL).
FT	TRANSMEM	278	301	7 (POTENTIAL).
FT	DOMAIN	302	352	CYTOPLASMIC (POTENTIAL).
FT	DISULFID	101	178	BY SIMILARITY.
FT	MOD_RES	3	3	SULFATION (BY SIMILARITY).
FT	MOD_RES	10	10	SULFATION (BY SIMILARITY).
FT	MOD_RES	14	14	SULFATION (BY SIMILARITY).
SQ	SEQUENCE	352 AA;	40445 MW;	4F8E4F344CEB7C91 CRC64;

Query Match 40.6%; Score 359; DB 1; Length 352;

Best Local Similarity 33.9%; Pred. No. 1.8e-27;

Matches 96; Conservative 8; Mismatches 29; Indels 150; Gaps 7;

Qy	20	MDYQVSSPIYDINYYTSEPCQKINVQIAA-----	49
		:	
Db	1	MDYQVSSPTYDIDYDTSEPCQKINVQIAARLLPPLYSLVFIFGFVGNMLVILVLINCKR	60
Qy	50	-----YKCGLC-----AAAQWDFGNTMCQHQRVHGHHHSYKCG---	83
		: :	
Db	61	LKSMTDIYLLNLAISDLFFLLTVPFWAHYAAAQWDFGNTMC--QLLTGLYFIGFFSGIFF	118
Qy	84	-----LCTRSQKEGLHYTC	97
		:	
Db	119	IILLTIDRYLAIVHAVFALKARTVTFGVTVSVITWVAVFASLPGIIFTRSQKEGLHYTC	178
Qy	98	SSHFPYSQYQFWKNFQTLKI-----HQRVHGG-----	124
		: :	
Db	179	SSHFPYSQYQFWKNFQTLKIVILGLVLPLLMVICYSGILKTLLRCRNEKKRHRVRLIF	238
Qy	125	-----GGSYKCGLC-----QEFGGLNNCSSSNRLDGHQRV	154
		:	
Db	239	TIMIVYFLEWAPYNIVLLLNTFQEFGGLNNCSSSNRLDQAMQV	281

Search completed: March 5, 2004, 16:23:45

Job time : 22.321 secs